

GEOLOGY

of

MC LEAN COUNTY, NORTH DAKOTA

by

John P. Bluemle

North Dakota Geological Survey

Grand Forks, North Dakota

1971

BULLETIN 60 – PART I
North Dakota Geological Survey
Edwin A. Noble, *State Geologist*

COUNTY GROUND WATER STUDIES 19 – PART I
North Dakota State Water Commission
Milo W. Hoisveen, *State Engineer*

Prepared by the North Dakota Geological Survey
in cooperation with the North Dakota State
Water Commission, the McLean County Board of Commissioners,
and the United States Geological Survey

GEOLOGY

of

MC LEAN COUNTY, NORTH DAKOTA

by

John P. Bluemle

North Dakota Geological Survey

Grand Forks, North Dakota

1971

BULLETIN 60 – PART I

North Dakota Geological Survey

Edwin A. Noble, State Geologist

COUNTY GROUND WATER STUDIES 19 – PART I

North Dakota State Water Commission

Milo W. Hoisveen, State Engineer

Prepared by the North Dakota Geological Survey
in cooperation with the North Dakota State
Water Commission, the McLean County Board of Commissioners,
and the United States Geological Survey

This is one of a series of county reports published cooperatively by the North Dakota Geological Survey and the North Dakota State Water Commission. The reports are in three parts; Part I describes the geology, Part II presents ground water basic data, and Part III describes the ground water resources. Parts II and III will be published later.

CONTENTS

| | Page |
|---|------|
| ABSTRACT | 1 |
| INTRODUCTION | 2 |
| Purpose | 2 |
| Methods of Study | 2 |
| Previous Work | 3 |
| Acknowledgments | 4 |
| Surface Topography and Physiography | 4 |
| STRATIGRAPHY | 6 |
| General Statement | 6 |
| Precambrian Rocks | 8 |
| Paleozoic Rocks | 8 |
| Sauk Sequence | 8 |
| Tippecanoe Sequence | 10 |
| Kaskaskia Sequence | 10 |
| Absaroka Sequence | 11 |
| Mesozoic and Tertiary Rocks | 11 |
| Zuni Sequence | 11 |
| Quaternary Rocks | 15 |
| Preglacial Sediments | 15 |
| Description | 15 |
| Origin | 15 |
| Coleharbor Formation | 16 |
| General Statement | 16 |
| Definition | 17 |
| Distribution | 21 |
| Boulder-clay | 21 |
| Sand and Gravel | 22 |
| Silt and Clay | 23 |
| Age and Origin | 23 |
| Postglacial Sediments | 24 |
| Alluvial Sediments | 24 |
| Slough Sediments | 25 |
| Eolian Sediments | 25 |
| Landforms | 25 |
| Erosional Forms | 25 |
| Depositional Forms | 26 |
| End Moraines | 26 |
| Ground Moraine and Sheet Moraine | 27 |

| | |
|---|-----------|
| Dead-Ice Moraine | 27 |
| Kettle Chains | 28 |
| Large Glacial Boulders | 29 |
| Glacio-Fluvial Forms | 30 |
| Alluvial Plains | 30 |
| Collapsed Alluvial Plains | 31 |
| Kames | 31 |
| Flat Lake Topography | 32 |
| Undulating Lake Topography | 32 |
| Meltwater Trenches | 32 |
| Erosion Surfaces | 33 |
| Terraces | 35 |
| Nonglacial Forms | 36 |
| River Floodplains | 36 |
| Sloughs | 37 |
| Dunes | 37 |
| Stream-Eroded Topography | 37 |
| GEOLOGIC HISTORY | 38 |
| Preglacial History | 38 |
| Topography on the Bedrock Surface | 39 |
| Development of Drainage During the Quaternary | 39 |
| Preglacial Drainage | 40 |
| Drainage Development During Glaciation | 40 |
| Glacial History | 43 |
| Introduction | 43 |
| Pre-Wisconsinan Drift | 43 |
| Dead Man Drift | 43 |
| Mercer Drift | 48 |
| Pre-Wisconsinan Glacial History | 52 |
| Early Wisconsinan Drift | 52 |
| Napoleon Drift | 52 |
| Early Wisconsinan Glacial History | 54 |
| Late Wisconsinan Drift | 55 |
| Lostwood Drift | 55 |
| Late Wisconsinan Glacial History | 58 |
| ECONOMIC GEOLOGY | 63 |
| Petroleum | 63 |
| Lignite | 63 |
| Clinker | 63 |
| Riprap and Building Stone | 63 |
| REFERENCES CITED | 64 |

ILLUSTRATIONS

| | Page |
|---|-------------|
| Plate | |
| 1. Geologic map of McLean County | (in pocket) |
| 2. Geologic sections of exposed bedrock in McLean County | (in pocket) |
| 3. Topographic map of the bedrock surface of McLean County | (in pocket) |
| Figure | |
| 1. Topographic and physiographic map of McLean County | 5 |
| 2. Regional map showing location of the Williston basin and major structural features | 7 |
| 3. Stratigraphic column of North Dakota | 9 |
| 4. Subcrop map of Cannonball, Tongue River and Sentinel Butte Formations in McLean County | 14 |
| 5. View across bay of Dead Man Coulee showing type area of the Coleharbor Formation | 19 |
| 6. Chunks of iron-cemented gravel conglomerate lying on boulder clay surface (the exhumed upper surface of the Dead Man till) | 19 |
| 7. Photo of type section of the Coleharbor Formation along Lake Sakakawea | 20 |
| 8. Esker between Benedict and Max in north-central McLean County | 29 |
| 9. Middle branch of Douglas Creek meltwater trench in T. 148 N., R. 85 W. | 34 |
| 10. Turtle Creek valley east of Washburn | 34 |

| | |
|--|----|
| 11. View south over the flat profile of a remnant of an erosion surface | 35 |
| 12. Map showing major preglacial drainages and glacial diversion trenches in the McLean County area | 41 |
| 13. Map of North Dakota showing maximum extent of pre-Wisconsinan, Early Wisconsinan and Late Wisconsinan glaciers | 44 |
| 14. Nearly vertical exposure of boulder-clay (Dead Man till) showing wave-weathered joint faces | 46 |
| 15. Inclusion of banded silts (lake sediments) in boulder-clay (Dead Man till) | 46 |
| 16. "Cobbles" composed of boulder-clay (Dead Man till) | 47 |
| 17. Caliche accumulations near the top of the Dead Man till | 47 |
| 18. Boulder-clay of unit 3 in description of type section of Coleharbor Formation (above shovel) | 48 |
| 19. Contact between till and fluvial facies of the Mercer drift | 49 |
| 20. Comparison of glacial stratigraphy in McLean County with that in the Red River Valley | 50 |
| 21. Map of McLean County showing areas of various drifts | 53 |
| 22. Receding early Wisconsinan Napoleon ice | 56 |
| 23. Deposition of the Blue Mountain end moraine | 59 |
| 24. Maximum extent of late Wisconsinan ice in eastern McLean County | 61 |
| 25. Withdrawal of the active ice margin from McLean County | 62 |

THE GEOLOGY OF MC LEAN COUNTY

by

John P. Bluemle

ABSTRACT

McLean County is in west-central North Dakota on the east side of the Williston basin. It is underlain by 8,500 to 13,000 feet of Paleozoic, Mesozoic, and Cenozoic rocks that dip to the west at low angles. The Tertiary, Cannonball, Tongue River, and Sentinel Butte Formations lie directly beneath glacial drift. The Tongue River and Sentinel Butte Formations are widely exposed in McLean County, and the Cannonball Formation is exposed in a few places. Glacial drift occurs throughout the county and reaches a maximum thickness of at least 400 feet in certain preglacial valleys. This glacial drift is assigned to a new geologic formation, the Coleharbor Formation, which is defined and described in this report.

The eastern part of McLean County is part of the Missouri Coteau and is characterized by a hilly surface on dead-ice moraine. Associated with the dead-ice moraine are numerous kames, lake plains, ice-contact gravel deposits, and areas of collapsed outwash topography. Much of the remainder of the county is part of the Coteau Slope that is characterized by a gently rolling surface on ground moraine. The Missouri Trench forms the western boundary of the county. It has steep slopes developed mainly on bedrock.

Pre-Wisconsinan glacial deposits were identified in McLean County but little is known of either their age or the circumstances under which they were deposited. Early Wisconsinan drift covers most of the area of the Coteau Slope and Late Wisconsinan drift covers most of the Missouri Coteau. Both the early and late Wisconsinan glaciers stagnated as they thinned and receded from the area resulting in hummocky areas of dead-ice moraine. The modern route of the Missouri River did not become firmly established until the late Wisconsinan glacier receded from the area.

INTRODUCTION

Purpose

This report is the first part in a three-part study of the geology and ground water resources of McLean County, North Dakota. It is being prepared cooperatively by the North Dakota Geological Survey, the North Dakota State Water Commission, the McLean County Board of Commissioners, and the U. S. Geological Survey. The remaining two parts of the study consist of a compilation of basic ground water data and a discussion of the ground water resources of the county.

The present report contains both description and interpretation of the geology of McLean County. Parts of the report that are primarily descriptive include the discussions of the topography, rock, and sediment of the county. This information is intended for use by anyone interested in the physical nature of the materials underlying the county. Such people may be water-well drillers or hydrologists interested in the distribution of sediments that might produce usable ground water, civil engineers and contractors interested in the gross characteristics of foundation materials at possible construction sites or the locations of possible sources of borrow material for concrete aggregate, industrial concerns looking for possible sources of economic minerals, residents interested in knowing more about their county, and geologists interested in the physical evidence for the geologic interpretations.

Parts of the report that are primarily interpretive include the discussions of the landforms and Quaternary history in McLean County. These parts are intended for those interested in the geologic processes and sequence of events during Pleistocene time.

Methods of Study

The writer mapped about three-fourths of McLean County during the 1966 and 1967 field seasons. The remaining one-fourth of the area was mapped during the 1966 field season by Jack Kume, formerly of the North Dakota Geological Survey.

Data were plotted on McLean County highway maps, scale 1:63,360, prepared by the North Dakota State Highway Department. In addition, topographic maps of the 7.5-minute series were available for much of the county and maps of the 15-minute series were available for most of the remainder of the area. Air photos, scale 1:20,000, taken

in 1958, were used to accurately place geologic contacts. The surficial mapping was done by driving along all section line roads and trails, and recording lithologies at all roadcuts or exposures. Less accessible areas were covered on foot. A shovel and soil auger were used to obtain lithologic information in areas of poor exposures. In addition, about 100 holes were bored by the North Dakota Geological Survey truck-mounted auger. This auger is capable of sampling to a maximum depth of 150 feet. The North Dakota State Water Commission provided a rotary drilling rig that was used during the 1967, 1969, and 1970 field seasons for about 40,000 feet of test drilling.

Previous Work

All or portions of McLean County have been included in several previous studies. In 1883, Chamberlain presented a map of the Missouri Coteau in "Terminal Moraines of the Second Glacial Epoch." Todd (1896) described the moraines of the Missouri Coteau. Leonard (1916) described the "pre-Wisconsin drift" of North Dakota. He placed the western limit of the Wisconsin drift at the front of the "Altamont Moraine." H. E. Simpson in 1942 described the geology of the Garrison Quadrangle in an unpublished Master's thesis at the University of Illinois. Townsend and Jenke (1951) named the "Max Moraine" for a part of what had previously been referred to as the Altamont Moraine. They described the Max Moraine as extending "from the vicinity of Bismarck, northwestward for 800 miles." Lemke (1960) briefly discussed the northeast corner of McLean County in a report on the Souris River area. In 1958, Lemke and Colton published a summary of the Pleistocene geology of North Dakota based on a study of large scale air photo stereopairs of the glaciated part of the state. In 1965, Lemke, Laird, Tipton, and Lindvall published a somewhat more comprehensive report on the Pleistocene geology of the state. Clayton (1966) added to and revised this. Also available is the *Preliminary Glacial Map of North Dakota* (Colton, Lemke, and Lindvall, 1963). Several geologic reports of the present county series are now available for the area near McLean County. They include Kidder County (Rau and others, 1962); and Burleigh County (Kume and Hansen, 1965). Field work has been completed by the North Dakota Geological Survey and the U. S. Geological Survey in several more nearby counties, and report preparation is now in progress. They include Mercer, Oliver, Sheridan, McHenry, Ward, Renville, and Mountrail Counties. In addition, several circulars describing samples from exploratory oil wells have been

published, and various general studies of North Dakota have included all or parts of McLean County.

Acknowledgments

The help of various individuals and agencies involved in this study is appreciated. Jack Kume, formerly of the North Dakota Geological Survey, mapped a large area of McLean County. Robert Klausung, author of parts II and III of this study, provided test hole data and other valuable information. I also wish to thank Lee Clayton of the Department of Geology at the University of North Dakota for reviewing this report. Roy Staiger, who operated the North Dakota Geological Survey auger, provided valuable test hole data on glacial drift thicknesses. Finally, I extend my appreciation to the landowners of McLean County who were very cooperative in providing data on private wells and allowing access to their property.

Surface Topography and Physiography

A topographic map of McLean County is shown on Figure 1. Four distinct physiographic areas occur in the county:

- 1) The extreme northeast corner of the county is part of the relatively level Drift Prairie that covers most of the northeast half of North Dakota. Drainage on this gently undulating glaciated plain is fair to good, although many shallow depressions retain water after heavy rains. Streams of the Drift Prairie of McLean County and nearby areas flow into the Sheyenne River, a tributary of the Red River of the North whose waters eventually reach Hudson Bay. Relief, which is generally less than 20 feet in a mile, is due almost entirely to irregularities on the surface of the ground moraine.

The edge of the Drift Prairie is marked by the Missouri Escarpment, a relatively smooth but stream-dissected slope that rises to the Missouri Coteau. Elevations rise from about 1,600 feet at the base of the escarpment to over 2,000 feet at the top, a change of about 400 feet in a distance of about 5 miles.

- 2) The Missouri Coteau, which lies at the top of the Missouri Escarpment, is an area of high local relief averaging about 30 to 35 feet between lows and adjacent highs. This hilly area extends from




FIGURE 1. Topographic and physiographic map of McLean County.

east-central South Dakota northwestward into western Saskatchewan. The Missouri Coteau covers approximately the northeast half of McLean County and accounts for about 40 percent of the total area of the county. A few short streams occur in the area but none transect it. The area is characterized by numerous sloughs, lakes, and closely-spaced hills. The entire width of the Missouri Coteau is the continental divide between Gulf of Mexico and Hudson Bay drainage.

3) The Coteau Slope lies southwest of the Missouri Coteau. Relief is moderate, generally less than 25 feet locally, but greater near some of the deeper valleys. These valleys carry small intermittent streams south and westward over the Coteau Slope to the Missouri River. Drainage is well-developed, and only a few lakes and sloughs occur in the area. The topography of the Coteau Slope is mainly stream-dissected bedrock with a veneer of glacial deposits. Elevations range between about 2,000 feet near the Missouri River to over 2,400 feet in the western part of the county.

4) The southwest boundary of McLean County is the Missouri River, which flows in the Missouri Trench. Above Garrison Dam, in western McLean County, the trench is generally 3 to 5 miles wide from rim to rim and has a maximum relief of 200 to 250 feet (lake level to trench rim). Below the dam it is 1 to 2 miles wide with a maximum relief of about 350 feet. The wider segment of the trench is flooded by Lake Sakakawea and has sides consisting mainly of badlands on the Tertiary Sentinel Butte Formation. There, discontinuous glacial drift occurs on the sides of the trench down to the level of the lake. Below the dam, the floor of the trench is a broad floodplain. Some badlands occur on the sides of the trench, but generally the sides are relatively smooth with terraces in some areas.

STRATIGRAPHY

General Statement

McLean County is situated on the east flank of the Williston basin, an intracratonic, structural basin containing a thick sequence of sedimentary rocks (fig. 2). The Williston basin had its beginning in Early Paleozoic time. Sedimentary rocks in western McLean County probably reach a total thickness greater than 13,000 feet, an estimate based on drill hole data and projection of rock units from wells in adjacent areas. No wells have yet been drilled that penetrated the entire





FIGURE 2. Regional map showing location of the Williston basin and major structural features. Lower part of illustration shows a generalized North Dakota stratigraphic column and cross section across the state. Modified from Carlson and Anderson, 1966, p. 1834.

sedimentary sequence. The rocks dip to the west at low angles in McLean County, from less than 25 feet per mile in the Upper Cretaceous to over 50 feet per mile in the Paleozoic rocks.

Precambrian Rocks

Precambrian rocks range in depth from about 8,500 feet in northeastern McLean County to more than 13,000 feet in the western part of the county. No wells have penetrated Precambrian rocks in McLean County and their composition is unknown.

Paleozoic Rocks

Paleozoic rocks range in thickness from about 4,000 feet in northeastern McLean County to over 8,000 feet at the western edge of the county. For purposes of discussion, the Paleozoic rocks can be subdivided into four sequences (fig. 3). A sequence is defined as the preserved sedimentary rock record bounded by major regional unconformities. The Paleozoic sequences recognized in McLean County are, in ascending order, the Sauk, Tippecanoe, Kaskaskia, and Absaroka, with the Absaroka Sequence extending to include Triassic rocks of the Mesozoic Era.

SAUK SEQUENCE

The Sauk Sequence is represented in the Williston basin by the Deadwood Formation of Late Cambrian to Early Ordovician age. The Deadwood is an onlap depositional sequence consisting primarily of a basal sandstone overlain by shale and carbonate and then by another sandstone. It thickens westward from a minimum of about 200 feet in northeast McLean County to a maximum of about 700 feet in the northwest part of the county. In the Vaughn-Hanson No. 1 well (Sec. 10, T. 146 N., R. 81 W.), 101 feet of limestone, shale and sandstone of the Deadwood Formation was penetrated. Fifty-eight feet of the Deadwood was penetrated in the Stanolind-McLean No. 1 well (Sec. 28, T. 150 N., R. 80 W.).

| SEQUENCE | SYSTEM | GROUP OR FORMATION | DOMINANT LITHOLOGY | |
|------------|---------------|--|---|------------------------------|
| TEJAS | | Colocharbor | Glacial Drift | |
| | | White River | Clay, Sand, Limestone | |
| ZUNI | TERTIARY | Golden Valley | Clay, Sand, Silt | |
| | | Fort Union Group | Sentinel Butte | Sandstone, Shale, Lignite |
| | | | Tongue River | Sandstone, Shale, Lignite |
| | | | Cannonball | Marine Sandstone, Shale |
| | | Hell Creek | Sandstone, Shale and Lignite | |
| | CRETACEOUS | Montana Group | Fox Hills | Marine Sandstone |
| | | | Pierre | Shale |
| | | | Niobrara | Shale, calcareous |
| | | Colorado Group | Carlile | Shale |
| | | | Greenhorn | Shale, calcareous |
| | | | Belle Fourche | Shale |
| | | Dakota Group | Mowry | Shale |
| | | | Newcastle | Sandstone |
| | | | Skull Creek | Shale |
| Fall River | | | Sandstone and Shale | |
| JURASSIC | Morrison | Lakota | Sandstone and Shale | |
| | | | Shale and Clay | |
| | Sundance | Shale, green & brown, Sandstone | | |
| | Piper | Limestone, Anhydrite, Salt and red shale | | |
| ABSAROKA | TRIASSIC | Spearfish | Siltstone, Salt and Sandstone | |
| | PERMIAN | Minnekahta | Limestone | |
| | | Opeche | Shale, Siltstone and Salt | |
| | PENNSYLVANIAN | Minnekahta | Sandstone and Dolomite | |
| | | Amaden | Limestone, dolomitic, Shale and Sandstone | |
| KASKASKIA | MISSISSIPPIAN | Big Snowy Group | Heath | Shale |
| | | | Otter | Sandstone |
| | | | Kibbey | Limestone |
| | | Madison | Limestones and Evaporites | |
| | DEVONIAN | Balkon | Siltstone and Shale | |
| | | Three Forks | Shale, Siltstone and Dolomite | |
| | | Birdbeak | Limestone | |
| | | Duperow | Dolomite and Limestone | |
| | | Souris River | Dolomite and Limestone | |
| | | Dawson Bay | Dolomite and Limestone | |
| | | Prairie | Halite | |
| | | Winnipegosis | Limestone and Dolomite | |
| | SILURIAN | Interlake | Dolomite | |
| TIPPECANOE | ORDOVICIAN | Stonewall | Dolomite and Limestone | |
| | | Stony Mountain Fm. | Gunston Mbr. | Dolomite and Limestone |
| | | | Stoughton Member | Argillaceous Limestone |
| | | Red River | Limestone and Dolomite | |
| | | Winnipeg Group | Roughlock | Calcareous Shale & Siltstone |
| | | | Icebox | Shale |
| | | | Black Island | Sandstone |
| SAUK | CAMBRIAN | Deadwood | Limestone, Shale and Sandstone | |
| | PRE-CAMBRIAN | | | |

FIGURE 3. Stratigraphic column of North Dakota. Heavy dashed lines represent major regional unconformities. Modified from Carlson and Anderson, 1966, p. 1835.

TIPPECANOE SEQUENCE

The Williston basin began to be a slightly negative area during deposition of the Tippecanoe Sequence. This sequence is the result of a transgressive event during which the seas invaded from the south and east, and the Williston basin became part of a much more extensive epicontinental sea. The Tippecanoe Sequence is represented in the area by rocks of Middle Ordovician to Silurian age. The initial deposits of the sequence were the clastics of the Winnipeg Group. These were followed by carbonates with minor amounts of evaporites of the Red River, Stony Mountain, Stonewall, and Interlake Formations.

In McLean County, the thickness of the Tippecanoe ranges from about 1,300 feet in the east to about 2,200 feet in the west. The two previously mentioned wells, the Vaughn-Hanson No. 1 and the Stanolind-McLean No. 1 are the only two that have penetrated the Tippecanoe Sequence in McLean County. In these wells, the thicknesses were 1,442 feet and 1,437 feet respectively.

KASKASKIA SEQUENCE

During deposition of the Kaskaskia Sequence, the Williston basin was slightly more tectonically negative than during the previous two sequences. The initial deposits of the Kaskaskia Sequence represent a transgressive sea that spread over the area from the north and west during Devonian time.

Devonian formations that have been recognized in the Williston basin include, in ascending order, the Winnipegosis (mainly carbonates), Prairie (mainly salt with some limestone and anhydrite), Dawson Bay (limestone and dolomitic limestone), Souris River (alternating limestone and thin argillaceous beds), Duperow (cyclical carbonates and shales), Birdbear (limestone), and Three Forks (shale, anhydrite, siltstone, and dolomite). The overlying Mississippian rocks were deposited mainly during normal marine conditions. They include rocks of the Bakken Formation (fine-grained clastics), the Madison Group (carbonates), and Big Snowy Group (shale, carbonates, and sandstones).

The Kaskaskia Sequence in McLean County ranges from about 2,000 feet thick in the southeast to about 4,100 feet thick in the northwest. In addition to the two previously mentioned wells, six others have penetrated the Kaskaskia Sequence to varying depths.

ABSAROKA SEQUENCE

The Absaroka sea spread across the area from the west. Deposition began with nonmarine mudstone, followed by relatively thin marine deposits and then predominantly clastic and evaporite redbed units. Rocks of the Pennsylvanian and Permian Systems and the Spearfish Formation of Permian-Triassic age comprise the Absaroka Sequence. Formations include, in ascending order, the Amsden (dolomite and limestone), Minnelusa (clastics), Opeche (redbed clastics), Minnekahta (dolomite and limestone), and Spearfish (redbed clastics). The Absaroka Sequence in McLean County ranges from about 250 feet thick in the northeast to about 1,100 feet thick in the southwest. Eight of the ten oil exploration tests drilled to date in McLean County have penetrated the Absaroka Sequence.

Mesozoic and Tertiary Rocks

Mesozoic rocks range in thickness from about 3,900 feet in northeastern McLean County to about 5,000 feet at the western edge of the county. All of these rocks are part of the Zuni Sequence with the exception of the previously discussed Spearfish Formation. The Tertiary rocks of McLean County range in thickness from about 400 feet in the northeast to more than 1,100 feet in the western part of the county.

ZUNI SEQUENCE

Mesozoic rocks of the Zuni Sequence in the Williston basin consist mainly of clastic rocks that were deposited in widespread Jurassic and Cretaceous seas. Jurassic strata range from about 600 to 1,000 feet thick in McLean County and consist of evaporites, red shale, and carbonates in the Piper Formation, and fine-grained clastics of the Sundance Group. Cretaceous rocks include well-developed sandstones in the Fall River-Lakota interval and a poorly developed sandstone in the Newcastle Formation. The rest of the Cretaceous rocks, below the Fox Hills Formation, are gray shales with some calcareous shales and some thin bentonites; they include the Skull Creek, Mowry, Belle Fourche, Greenhorn, Carlile, Niobrara, and Pierre Formations.

The Fox Hills Formation conformably and gradationally overlies the Pierre Formation. It is a marine sandstone and shale sequence that

ranges in thickness from about 350 to 450 feet in McLean County with the greatest thicknesses in the northeastern part of the county. The Hell Creek Formation, the youngest Cretaceous formation, conformably overlies the Fox Hills Formation in McLean County. It is of continental origin and consists of interbedded gray, greenish-gray and brown sandstones, mudstones, siltstones, carbonaceous shales, and thin lignite seams. It is overlain by the interbedded sandstones, shales, and lignite beds of the Paleocene age Ludlow Formation. The Hell Creek and Ludlow Formations have a combined thickness of about 100 feet in eastern McLean County to about 500 feet in the western part of the county.

The remaining Tertiary rocks of McLean County, in ascending order, are the Cannonball, Tongue River, and Sentinel Butte Formations. These sediments are all of the Zuni Sequence and late Paleocene in age. They are represented in the Williston basin by predominantly non-marine deposits derived from source areas to the west.

The Cannonball Formation, which conformably overlies the Ludlow Formation, is probably a marine equivalent to the Ludlow. It consists of olive black, carbonaceous and lignitic siltstone and shale, lignite, and micaceous friable sandstones. They grade upward into similarly colored noncalcareous siltstones and claystones and friable, glauconitic, noncalcareous sandstones.

The Cannonball Formation in the western and southern part of McLean County is as much as 300 feet thick. It thins northeastward to less than 100 feet thick. Exposures of the uppermost Cannonball beds occur in Sections 22 and 23, T. 143 N., R. 81 W. in southern McLean County. The contact of the highly weathered, brownish-black Cannonball shales with the overlying Tongue River sandstones occurs at an elevation of about 1,680 feet in this area, a figure that seems anomalously low in view of the fact that it is at least 1,700 feet across the river in Oliver County in a regionally downdip direction. Some minor structure may be indicated in the area.

The Tongue River and Sentinel Butte Formations, continental formations of Paleocene age, are the youngest bedrock formations in the county. They directly underlie the glacial drift, except in a small area in extreme southern McLean County where they are absent, and the Cannonball Formation underlies the drift, and in northwestern McLean County where preglacial Pleistocene sands lie between them and the overlying drift. Where the drift is thin or absent they are exposed; the best and most extensive exposures occur in and near the Missouri Trench, where headward erosion has formed areas of badlands topography. Maximum thickness of the two formations is about 800 feet in western McLean County.

Exposures of the Tongue River Formation, which underlie the Sentinel Butte Formation, are confined to approximately the eastern third of the county; the Sentinel Butte occurs over the remainder of the county (fig. 4). The contact between the two formations is difficult to recognize in McLean County. Based on tentative interpretations of several measured sections, the contact is exposed in the vicinity of Garrison Dam at an elevation of about 1,900 feet (Plate 1). The contact between the Tongue River and Cannonball Formations in this area is at about 1,500 feet, based on test hole data to the southwest in Mercer County; therefore, the Tongue River is about 400 feet thick in the area. However, about 450 feet of composite section was measured in the area. This footage includes about 120 feet of section that was poorly exposed due to slumping, so the possibility for error is considerable.

The Tongue River consists of buff to orange buff and gray sands, silts, and clays that range from poorly to fairly well cemented. The sands are commonly cross-bedded and channeled in places and have local horizons of sandstone concretions. Some of the more indurated sand horizons form prominent ledges that can be carried for several hundred yards. Fossils collected from the Tongue River Formation include mainly gastropods with some pelecypods and ostracods.

The clays and silts of the Tongue River form good marker beds for correlative purposes. They are commonly light gray with orange bands due to iron oxide-rich beds. Commonly associated with the clay and silt beds are lignitic zones and lignite beds. Most of these beds are less than a foot thick, and no commercially useful lignite was found in the Tongue River Formation of McLean County.

About 200 feet of Sentinel Butte section was measured in western McLean County. The lowermost part of the formation is too poorly exposed in the area to allow for accurate correlation with bedrock exposures near Garrison Dam. In general, the Sentinel Butte in McLean County consists of light gray to brownish gray sands and silts. Petrified wood, mainly *Metasequoia*, is common in the Sentinel Butte Formation, although some occurs in the Tongue River as well. In some badland areas, large stumps and logs occur and small chips of petrified wood cover the surface. Reddish scoria beds, formed when burning lignite beds baked surrounding sediments, are common in both the Tongue River and Sentinel Butte Formations. Their lateral continuity is restricted to less than a few miles so they can be used only for very local correlations. Large, tubular concretions are common in the Sentinel Butte Formation. They are round to oval in cross-section; often they stand on pedestals. They consist of cross-bedded sands that are commonly coarser than the surrounding sediments.

Several beds of lignite up to 2 or 3 feet thick occur, and two beds are being mined. One of these mines, the Underwood Coal Company




FIGURE 4. Subcrop map of Cannonball, Tongue River and Sentinel Butte Formations in McLean County. These three bedrock formations lie directly beneath the glacial drift except in areas where glacial deposits are absent and in a small area of the northwestern corner of the county where preglacial Quaternary (?) river sediments occur between the bedrock and the drift.

Mine, is in Section 23, T. 146 N., R. 82 W. Here, an 8- to 10-foot thick bed is being mined. This lignite occurs at an elevation of 1,950 to 1,960 feet, about 25 feet above the presumed base of the Sentinel Butte Formation in that area. The other mine is the B & W Coal Mine (Sec. 16, T. 148 N., R. 86 W.) where two lignite beds are being mined. The lower 5-foot-thick bed is separated from the upper 2-foot-thick bed by a 4-inch clay seam. The lignite occurs at an elevation of about 1,915 to 1,925 feet, about 160 feet above the base of the Sentinel Butte Formation. Both of these mines are relatively small operations.

The single most important criterion for differentiating the Tongue River and Sentinel Butte Formations is color. In western North Dakota, oxidized exposures of the Tongue River have light, buff-yellow colors with considerable variations in texture and color between individual beds and, particularly when viewed from a distance, they present a bright overall aspect. In contrast, the Sentinel Butte beds are more gray and somber in color and rather uniform in both texture and color.

Quaternary Rocks PREGLACIAL SEDIMENTS

Description

In several test holes in T. 150 N., R. 88 W., in extreme northwestern McLean County, fluvial sediments consisting of silts, sands, and gravels were found beneath glacial deposits. The trend of the sediments could not be determined. They do not appear to follow a valley as they are not particularly deep. The gravels and sands consist mainly of coarse, dark brown cherts, limestones, dolomite, and fragments of lignite. Silts and sands overlies the gravels and these are in turn overlain by clays and silts.

Origin

The lithologies of the fluvial sediments described above suggests that they were derived mainly from the Rocky Mountains or Black Hills and that they were deposited by rivers flowing northeastward in late Tertiary or early Pleistocene time. The sediments follow no particular trend, and if they were deposited in a valley, that valley has been destroyed and may even be a topographic high today. Such a situation implies a long interval of time between deposition of the sediments and overriding by glaciers. During this time, erosion over the area completely changed drainage patterns so that there now is no evidence for any buried valley. For this reason, it seems likely the sediments are considerably older than fluvial sediments that are found in buried

valleys, perhaps even pre-Pleistocene in age. The presence of what are apparently lake deposits of Pleistocene age on top of the fluvial sediments suggests that a lake formed when early Pleistocene ice blocked the drainage.

Stratigraphic situations similar to the one described above occur elsewhere in McLean County, but these are mainly in northeast-trending preglacial valleys (fig. 12), and in most cases they are documented only by single test holes. Preglacial fluvial sediments may underlie the glacial deposits in many of the preglacial valleys.

COLEHARBOR FORMATION

General Statement

Several surface glacial drift units have been recognized and named in North Dakota. These are inferential units recognized on the basis of ecologic analysis, interpretation of mineralogy, depositional structures and textures, and fossil faunas. In North Dakota, most of the named drift units are thought to have been deposited during single phases of activity of the glacier, and they can be related to some event in the history of the glacier. Because the drift units are recognized mainly on the basis of inferred geologic history or origin, they are highly interpretive and subject to change as more knowledge becomes available.

The surface drifts that have been named in North Dakota are, for the most part, lithologically indistinguishable on a regional scale. For this reason, and because of their inferential nature, it is practical to include all of them in a single lithostratigraphic unit. Lithostratigraphic units, or rock units, are bodies of strata identified by objective lithologic criteria. They are delineated in vertical succession by surfaces representing changes in lithologic character or breaks in the depositional continuity (Krumbein and Sloss, 1963, p. 332). In North Dakota, all the glacial sediments belong to the Coleharbor Formation, which is one of the most widespread geologic units in the state.

The glacial sediments in North Dakota have been studied sufficiently to adequately characterize and formally designate the Coleharbor Formation entirely in lithostratigraphic terms without resort to inferred geologic history or origin. The Coleharbor Formation consists of thousands of alternating beds but only three main facies: 1) interlayered bouldery, cobbly, pebbly, sandy, silty clay; 2) sand and gravel; and 3) silt and clay. These three facies taken as a whole form as uniform and distinctive a unit as any of the previously named formations in North Dakota. The Coleharbor Formation ranges from 0 to 600 feet thick in North Dakota.

By designating a formal rock unit, the Coleharbor Formation, it becomes possible to avoid the common practice of treating glacial deposits and bedrock in two entirely different ways. The formal designation should also facilitate more convenient study of comparable rocks over wide areas, not only in North Dakota, but in surrounding areas as well.

Christiansen (1967, 1968) has recognized several different units of glacial drift in the Saskatoon, Saskatchewan, area. He has divided the drift, first of all, into the Sutherland and Saskatoon Groups on the basis of carbonate content of tills, weathering zones, and intertill stratified drift. He has further subdivided the upper group, the Saskatoon, into the Battleford and Floral Formations.

Subdividing the Coleharbor Formation in North Dakota on a lithostratigraphic basis is not yet practical. Preliminary field work with recognized drift units in North Dakota (see the discussion of the Quaternary History later in this report) suggests that Christiansen's Battleford Formation is equivalent to the Lostwood drift and perhaps to the Napoleon drift in North Dakota. His Floral Formation may be equivalent to the Mercer and Dead Man drifts. Further study may make it possible to subdivide the Coleharbor Formation into members, or it may become necessary to elevate the Coleharbor Formation to group status. Possible members may be found to coincide with different glacial stages, or they may reflect predictable and mappable differences in till carbonate content, degree of weathering of feldspars, clay content, or some as yet unknown physical characteristics.

Definition

The term "Coleharbor Formation" is proposed for the lithostratigraphic unit that includes all bouldery, cobbly, pebbly, sandy, silty clay, sand and gravel, and silt and clay exposed in the type section and in all comparable sections. It is named for the town of Coleharbor in McLean County, North Dakota. The best exposures of the formation in North Dakota are located about 6 miles west of Coleharbor along the shore of Lake Sakakawea in Secs. 14, 15, and 22, T. 147 N., R. 84 W. ($101^{\circ} 21' W.$ Longitude, $47^{\circ} 32' N.$ Latitude); this will be considered the type area (fig. 5). Here, rapid erosion of the shore of the lake by wave action has exposed as much as 65 feet of nearly vertical section. The type section of the Coleharbor Formation follows. It is located about 300 yards south of the mouth of Dead Man Coulee in the NW $\frac{1}{4}$, NW $\frac{1}{4}$, NE $\frac{1}{4}$, Sec. 22, T. 147 N., R. 84 W., McLean County, North Dakota (fig. 7).

8. *Boulder-Clay*, 15 to 20 feet thick; brownish buff to gray, rather sandy with a few granitic and carbonate pebbles; chunky, unjointed and vaguely layered; calcareous. ORIGIN: Napoleon till of early Wisconsinan (?) age.
7. *Gravel*, 0 to 5 feet thick; medium-grained, sandy, cross-bedded in places; some iron staining near base of unit; consists mainly of igneous and carbonate pebbles. ORIGIN: Stream channel deposit.
6. *Boulder-clay*, 25 feet thick; brownish gray, sandy and pebbly with a few lignite chips and pockets of reddish brown, finely divided limonite; strong vertical jointing results in almost vertical cliffs; appears to have good vertical permeability. ORIGIN: Mercer till of pre-Wisconsinan age.
5. *Sand*, 2 to 4 feet thick; fine to medium grained, cross-bedded and loose; consists mainly of quartz with much finely-divided lignite; lies above unit 4 in most places but appears to be laterally equivalent to unit 4 in a few places. ORIGIN: Glacial outwash deposit.
4. *Gravel*, 2 to 3 feet thick; coarse, sandy, highly iron-cemented into a conglomerate (fig. 6); chunks up to 10 feet long, but averaging 3 to 4 feet long, have fallen intact into the reservoir; consists mainly of pebbles and cobbles with lithologies typical of glacially-derived gravels from the northeast in Canada (igneous and carbonate rocks). ORIGIN: Glacial outwash deposit.
3. *Boulder-clay*, 15 to 20 feet thick; mottled gray and yellowish buff with lighter shades overall than found in overlying boulder-clay units; cliff surfaces are oxidized, but in freshest cuts the unit is unoxidized at 10 to 15 feet below its upper surface; silty, very pebbly with sand pockets up to 6 inches across, large inclusions of lignite and lake sediments, and a few large boulders up to 15 feet across; abundant lignite chips and charcoaled wood fragments; relatively more carbonate pebbles than in overlying boulder-clay units; irregularly jointed with selenite crystals, iron oxide and manganese dioxide staining common on joint faces; partially lithified (breaks across pebbles in places) and massive; a few iron concretions; upper contact of this unit with the overlying gravel is very sharp and, in places where gravel has been washed off by waves, a relatively flat erosion surface has been exhumed. ORIGIN: Dead Man till of pre-Wisconsinan age.




FIGURE 5. View across bay of Dead Man Coulee showing type area of the Coleharbor Formation. Cliffs in the distance are about 40 to 50 feet high.



FIGURE 6. Chunks of iron-cemented gravel conglomerate lying on boulder-clay surface (the exhumed upper surface of the Dead Man till). Photo taken near the area of the type section.





FIGURE 7. Photo of type section of Coleharbor Formation along Lake Sakakawea. Total exposed section shown here is about 40 feet. Numbers on diagram identify lithologic units on photo and correspond to numbers used in section description.

2. *Silt*, 3 feet thick (at water level and poorly exposed); brownish to yellowish gray; clayey; good horizontal banding that in places has been highly contorted; appears to be laterally equivalent to unit 1 in places. ORIGIN: Overridden proglacial lake deposit.
1. *Gravel*, 2 to 3 feet thick (poorly exposed at water level); some iron stains; mineralogy of pebbles and cobbles is typical glacially-derived assemblage (carbonates, granitics, and metamorphics). ORIGIN: Glacial outwash deposit.

Distribution

The Coleharbor Formation covers most of the northeastern two-thirds of North Dakota, eastern South Dakota, Saskatchewan, Manitoba, northern Montana, and western and southern Minnesota. The southern limit of the Coleharbor Formation is the limit of glaciation in South Dakota. Its eastern limit is probably in central Minnesota where reddish-yellow, sandy, noncalcareous and nonmontmorillonitic drift replaces it. The northeastern limit is probably near Lake Manitoba where sandy, highly calcareous nonmontmorillonitic drift replaces it. The northern and western limits are probably near the Canadian Shield and Rocky Mountains. It unconformably overlies older rocks of all ages and is overlain in many areas by sediments of Holocene age. The Coleharbor Formation is largely confined to areas of clay-rich, montmorillonitic Cretaceous and early Tertiary formations of the northern Great Plains.

The formation underlies about 95 percent of McLean County (pl. 1). It is more than 400 feet thick in at least one buried valley in western McLean County (Sec. 32, T. 149 N., R. 87 W.) but it averages about 120 feet thick throughout the county. Near the Missouri River trench and over much of western McLean County, the formation averages less than 50 feet thick.

Boulder-clay

The Coleharbor Formation of McLean County consists of about 70 percent bouldery, cobbly, pebbly, sandy, silty clay (boulder-clay for short) by volume. About 75 percent of the surface area of the county is boulder-clay but much of this area is underlain by thin deposits; the areas of sand and gravel are generally underlain by thicker deposits. The boulder-clay facies of the Coleharbor Formation is a relatively uniform, nonbedded mixture of approximately equal parts of sand, silt, and clay sized materials along with small percentages of pebbles, cobbles, and boulders as much as a few feet in diameter.

The boulder-clay is generally dark gray below the ground-water table, grayish brown above the water table. Colors generally become

darker downward from more to less highly oxidized zones. Boulder-clay is relatively more uniform mineralogically than most other sediment in the county. Montmorillonite and other clay minerals along with small amounts of carbonates, quartz, and feldspars comprise the clay-sized fraction of the boulder-clay. The silt and sand fraction consists largely of quartz and feldspar with some carbonates, shale, and lignite. Pebbles are varied and consist of limestone, dolomite, various hard metamorphic and igneous rock types, shale, and lignite. Most of the cobbles and boulders are igneous and metamorphic rock types; a few are carbonates.

Based on 78 analyzed samples from depths of 0 to 100 feet, the boulder-clay has a natural dry unit weight ranging from 92 to 122 pounds per cubic foot with an average of 111, a natural moisture content ranging from 16 to 28 percent with an average of 20.8 percent, and a triaxial compressive strength ranging from 17 to 223 pounds per square inch with an average of 54. Compressive strengths tend to increase with depth.

Because of its low permeability, the boulder-clay is generally not a satisfactory source of ground water. Any permeability that the boulder-clay has is due to vertical fracturing so that the permeability is greater vertically than horizontally.

Sand and Gravel

In McLean County, the Coleharbor Formation consists of about 25 percent sand and gravel by volume. About 20 percent of the surface area of the county is sand and gravel. The deposits occur both as thin layers and lenses within the boulder-clay and as thick, continuous sequences independent of the boulder-clay. The sand and gravel facies is generally horizontally layered except in the hilly areas in the eastern part of the county where the layering is commonly tilted, contorted, or faulted. Quality ranges from sandy gravel and gravelly sand that is relatively free of finer materials to very dirty gravel with high percentages of silt and clay included.

The sand and gravel facies is more variable in grain-size composition than is the boulder-clay facies. In general, the best sorted deposits underlie the flat expanse of land northwest and east of Turtle Lake and east of Washburn. These commonly have maximum grain sizes of less than 2 inches. Well-sorted sands and gravels occur also on valley terraces south of Ruso in northeastern McLean County. Much of the sand and gravel that occurs in irregularly-shaped ridges and mounds in the hilly areas of eastern McLean County is poorly sorted and coarse-grained. It contains considerable amounts of boulder-clay and boulders of all sizes and is generally of little or no economic value. The areas of sand and gravel deposits are shown on Plate 1 but no quality

distinction is made. If sand or gravel of a particular grain size is wanted, the areas shown should be sampled to determine favorable locations.

The sand and gravel facies of the Coleharbor Formation has a mineralogic composition similar to the boulder-clay. The mineralogy indicates that it was ultimately derived from the northeast in Canada. The sand-sized fraction is largely quartz and feldspars with minor amounts of shale and carbonates. The gravel-sized fraction is largely granitic and metamorphic rock types, limestone, and dolomite. Some of the gravel has a high percentage of shale. Near the surface, caliche (CaCO_3) coats the undersides of pebbles and cobbles.

Generally the sand and gravel is loose, but in random local occurrences it is cemented with iron-oxide and forms a conglomerate. Exposures of iron-cemented gravels are particularly common along the shore of Lake Sakakawea at the mouths of Snake Creek and Dead Man Coulee (T. 147 N., R. 84 W.). In areas where the sand and gravel is loose and uncemented, it is also highly permeable. The sand and gravel is generally the largest and most dependable source of high-quality groundwater in the county.

Silt and Clay

Approximately 5 percent of the Coleharbor Formation in McLean County consists of layers and lenses of silt and clay. Only about 2 percent of the surface area is silt and clay, but considerable thicknesses of it occur in buried valleys throughout the county. It commonly has horizontal layering a fraction of an inch in thickness. Sand is uncommon and there are very few pebbles.

The silt and clay facies has the same mineralogy as the silt and clay fraction of the boulder-clay: montmorillonite and other clay minerals, carbonates, quartz, and feldspars. Its permeability is about the same as the boulder-clay.

Where it is exposed at the surface in McLean County, the silt-sized fraction is commonly prevalent over the clay, but there are numerous exceptions. The same is true to a lesser extent of the subsurface deposits. The silt and clay deposits are not at all uniform throughout the county.

Age and Origin

The Coleharbor Formation was deposited during the ice age from several hundred thousand to about 9,000 years ago. The boulder-clay facies of the formation is mainly glacial till consisting of material eroded by the glaciers from areas north of McLean County as well as locally. Part of the boulder-clay was subsequently deposited directly from the moving ice, and the remainder slid, slumped, and flowed to its present position when the ice finally melted.

The sand and gravel of the Coleharbor Formation was deposited mainly by rivers and streams during glacial times. Some of these rivers and streams must have been fed by meltwater from the glaciers, but probably large amounts of sand and gravel were deposited by nonglacial rivers 12,000 to 9,000 years ago when precipitation amounts were high and runoff was much greater than it is today. Some of the sand and gravel was deposited on beaches of lakes.

The Coleharbor silt and clay exposed at the surface in eastern McLean County was deposited in lakes that were at least in part enclosed by glacial ice. Part of the surface silt and clay in western McLean County may have formed in the same way, but the remainder probably was deposited in natural lakes (not ice-walled) that would be in existence today if precipitation were greater. The subsurface silt and clay that is confined to buried valleys was probably deposited in large lakes that formed when easterly-flowing rivers were dammed by the advancing ice.

POSTGLACIAL SEDIMENTS

Holocene sediments have been deposited since the ice age throughout McLean County, especially beneath river and stream valley floodplains (pl. 1), beneath slough floors, and in sporadic occurrences of sand dunes.

Alluvial Sediments

Alluvial sediment occurs along stream and river channels throughout McLean County. It ranges from a few feet thick along some of the smaller streams to over 100 feet thick in places beneath the Missouri River floodplain. It is sometimes difficult to distinguish modern alluvial deposits from the underlying fluvial and lacustrine sediments of glacial age.

The alluvial sediments consist mainly of clay, silt, sand, and gravel. Silts and clays commonly comprise about 90 percent of the sediments, sand and gravel the remaining 10 percent. Distinct and separate horizontal beds of dark brown sand, silt, and clay are common in the alluvium. The alluvial silt and clay differs from silt and clay of the Coleharbor Formation in that it is commonly sandy with a high organic content in places. The sand and gravel is similar to that in the Coleharbor Formation, but it is much less extensive and occurs as small lenses within the silt and clay.

Slough Sediments

The sediment in the bottoms of sloughs is a few tens of feet thick and consists of very dark brown and black clays with a high organic content. Most of the slough sediments are located in the hilly eastern and northeastern parts of McLean County. They consist of material washed into the lower areas from adjacent hill-slopes by runoff water. Individual deposits are too small to map, and although many hundreds of sloughs containing clays exist, they are not shown on Plate 1.

Eolian Sediments

Deposits of silt a few feet thick are common on and adjacent to sand and gravel deposits. They were deposited during dry periods throughout the 9,000 years since the close of the ice age. The silt is generally of uniform texture, massive and blocky in cuts so that it stands in vertical exposures. Most exposures are buff colored. Windblown silt deposits occur in southern McLean County in Tps. 143 and 144 N., Rs. 80 and 81 W., and Tps. 146 and 147 N., Rs. 83 and 84 W., near Garrison Dam. The deposits in southern McLean County were not mapped as they are very discontinuous. The silt deposits near the Garrison Dam, which are about 15 to 20 feet thick, are well-exposed in the steep bluffs north of the dam. At least two paleosols occur in the loess in this area. The lower one is very continuous and is truncated by modern valleys, so it must be older than are the valleys. The upper one, which becomes multiple in a few places, follows modern surface topography.

Windblown sand deposits occur throughout McLean County; but, with one exception, they are small and insignificant. The exception is in Sec. 13, T. 145 N., R. 84 W., and Sec. 18, T. 145 N., R. 83 W., where large sand dunes occur. This deposit averages about 20 feet thick and consists mainly of brownish, fine-grained sand, mainly quartz. A few cobbles can be found in the area, but it seems likely they were carried there, probably by Indians, for anchoring tepees.

Landforms

EROSIONAL FORMS

Although erosional forms are common in some areas, only a few such landforms were recognized in McLean County. Small drumlin-like ridges that occur in extreme northeastern McLean County apparently formed during the last southeastward movement of ice northeast of the Missouri Coteau. The ridges are of erosional and/or depositional origin and were formed when ice moving over the fluvial deposits in the area

reshaped the gravels into linear forms. The ridges are composed of the same gravel as the underlying fluvial deposits plus some clay and silt. Cuts through the ridges reveal a till-like fabric in places.

DEPOSITIONAL FORMS

Nearly all the landforms in McLean County that were formed as a direct result of glacial action are of a depositional nature. Included in this category are the various types of moraine, regardless of composition, although it should be recognized that fluvial and colluvial processes have played an important role in determining the final characteristics of the features.

End Moraines

An end moraine is a linear deposit of glacial drift formed at or near the terminus of an ice sheet during a standstill of the glacier terminus. Most end moraines result from a process of dumping of superglacial till that has been brought into the ice by shearing at the edge of the glacier and then concentrated on the surface by ablation. A process of ice shove, similar to a bulldozing effect at the terminus, has been suggested as an important mechanism in the formation of some end moraines. Another type of end moraine, the shear moraine, is a special type of dump moraine. This is a smaller, isolated ridge of till that is seemingly unrelated to any major standstill of the glacier. It is produced by shearing either at the zone separating the active ice from the inactive ice, or at sites of bedrock highs. Many washboard moraines probably form in this way.

The most important characteristic of an end moraine is its linearity, either an overall linearity of the end moraine as a unit, or linearity expressed by the alignment of depressions and ridges within the moraine. The glacial drift is normally significantly thicker beneath the end moraine than on the adjacent ground moraine, although this is not a necessary requirement if it can still be demonstrated that the end moraine does mark an ice marginal position that persisted for some time. End moraines on the Missouri Coteau are best distinguished from the surrounding dead-ice moraine by the presence of these lineations parallel to the long axis of the till ridge. Commonly, the boundaries between dead-ice moraine and end moraine are obscured by the collapse and obliteration of the linear ridges. In McLean County, no end moraines were recognized on the Missouri Coteau.

The Blue Mountain end moraine occurs on the Coteau Slope of western McLean County. It has a maximum local relief of about 60 feet and elevations ranging between 2,000 and 2,200 feet. In addition to

numerous small ridges that parallel the overall northeast-southwest trend of the feature, abundant small disintegration features more characteristic of dead-ice moraine occur over most of the Blue Mountain end moraine. The glacial drift beneath the end moraine averages over 60 feet thick based on about a dozen test holes. This compares with an average thickness of about 25 feet for the drift ahead of the end moraine and less than 20 feet for the drift behind it, although the ground moraine behind the end moraine becomes thicker further northwest. Areas of high bedrock associated with the Blue Mountain end moraine, particularly in front of it, appear to have been important in determining the position of the end moraine.

Ground Moraine and Sheet Moraine

Much of the western and southern parts of McLean County are covered by ground moraine and sheet moraine (pl. 1). Areas of sheet moraine are characterized by a thin, discontinuous layer of glacial drift, chiefly till, draped over stream-eroded topography. Sheet moraine is a mappable, blanket-like, veneer of drift that was deposited directly from glacial ice (Kume, 1965). It has low glacial constructional relief with few kettles. Its overall shape is the same as that of the landscape over which it is draped; it has inherited a pre-existing shape. Sheet moraine forms when the glacier, which is thin and probably relatively clean, drops its load as it loses the competency to carry it.

In McLean County, most of the sheet moraine consists of early Wisconsinan Napoleon drift. It is discontinuous near the Missouri River trench of western McLean County, where headward erosion is removing it, and near some of the larger streams.

Ground moraine is a landform composed of a drift accumulation, chiefly till, that was deposited directly from a moving glacier behind its margin. It has relatively low constructional relief. The ground moraine of McLean County occurs mainly northwest of the Blue Mountain end moraine and was deposited by the glacier that deposited the end moraine. It is characterized by irregular to gently rolling topography with relief that averages less than 20 feet in a mile except near streams where it is greater.

Dead-Ice Moraine

Most of the dead-ice moraine in McLean County occurs on the Missouri Coteau (fig. 1). However, another band of relatively subdued dead-ice moraine occurs in the Underwood area to the west of the Coteau. The area of dead-ice moraine is characterized by knob and kettle topography, non-integrated drainage with numerous small ponds and sloughs and a bouldery surface on till. The relief of the dead-ice moraine in eastern and northern McLean County ranges up to 100 feet

locally, but it is only about 20 to 25 feet in a mile on dead-ice moraine in the Underwood area (pl. 1). Relief may differ greatly within short distances, and boundaries of differing degrees of relief are difficult to delineate. Areas of high relief are differentiated from areas of low relief on Plate 1.

Disintegration ridges are, perhaps, the most diagnostic characteristic of the dead-ice moraine. They are rectilinear or circular ridges of glacial drift that were deposited between, around, or within depressions in blocks of stagnant glacier ice, through the process of mass movement from above and/or squeezing from below. Such features are abundant on the dead-ice moraine of eastern McLean County.

Closed disintegration ridges are the most common type of disintegration ridge found in McLean County. They are not easily seen in the field, but are obvious on air photos. They average about 500 feet in diameter, but are as large as 1/2 mile, average about 15 feet in height and have circular to irregular shapes. Many of the closed disintegration ridges are breached at both ends, forming features that are easily recognized on air photos.

Except for a few of the larger ones, which are shown on Plate 1, the disintegration ridges were not differentiated. The ridges shown on Plate 1 are mainly eskers, a particular type of disintegration ridge that was deposited by water flowing in valleys on the stagnant glacial landscape. When the stagnant ice melted, the resulting gravel deposits remained as esker ridges. The largest esker in McLean County (fig. 8) occurs in T. 150 N., R. 82 W., and extends from Section 7 to 35, a distance of about 6 miles. It has two branches on the north and grades into collapsed outwash on the south. The esker rises a maximum of about 50 feet above the surrounding dead-ice moraine. Notable esker ridges also occur in T. 147 N., R. 80 W., northeast of Turtle Lake, and in T. 147 N., R. 79 W.

Kettle Chains

Most of the kettle chains in McLean County occur in areas of dead-ice moraine (pl. 1). These may be defined simply as areas with high concentrations of kettles or potholes that follow a linear pattern. Many of the kettles contain sloughs or lakes. The kettles commonly follow a poorly defined valley or otherwise topographically low area. Some kettle chains overlie valleys that became filled with glacial ice, and others may simply represent areas where particularly large pieces of ice became incorporated in the drift. The longer kettle chains probably have the best chance of overlying gravel-filled valleys, and test drilling for water might be most useful in these areas.



FIGURE 8. Esker between Benedict and Max in north-central McLean County (Sec. 21, T. 150 N., R. 82 W.). Relief on this esker is about 50 feet. Numerous small pits in the esker exposed generally coarse, poorly sorted gravel and sand.

Large Glacial Boulders

Dogden Butte in northeastern McLean County (T. 150 N., R. 79 W.) is apparently a large glacial erratic. North Dakota State Water Commission test hole 4079 shows a sequence of 100 feet of glacial till and fluvial deposits overlying 212 feet of bedrock (Tongue River), which in turn overlies 44 feet of till and fluvial deposits. Similar large glacial erratics composed mainly of bedrock have been observed by the writer in Ransom and Nelson Counties, North Dakota. To the east in Sheridan County, several dozen similar abrupt and prominent hills were mapped by the writer in 1969 (Bluemle, 1970). In that area, a depression is located "up-ice" from each hill. Although no obvious depression is associated with Dogden Butte, it seems likely the butte formed in the same way as did the Sheridan County hills.

In the Sheridan County area, which was also near the Missouri Escarpment, it was concluded that normal local groundwater flow systems interacted with the groundwater flow systems introduced by

the glacier, resulting in greatly increased groundwater discharge and accompanying high hydrodynamic pressures in the subsurface between the glacier and the Missouri Escarpment. The high pressures forced large blocks of materials, including bedrock and glacial drift, upward into the advancing glacier. These materials were subsequently deposited by the glacier on top of the Missouri Coteau, resulting in high relief dead-ice moraine that developed when large-scale glacial stagnation occurred. When the glacier receded from the area, hydraulic conditions were essentially identical to those that had existed when the glacier advanced. Because the ice margin was fluctuating, but generally receding, newly formed blocks of material were moved only short distances, and the adjacent depressions were left essentially undisturbed.

A depression may have existed north of Dogden Butte off the coteau, but that area was overridden by ice from the northwest after Dogden Butte is thought to have formed. This later advance probably filled any existing depression with drift.

GLACIO-FLUVIAL FORMS

Streams flowing on or from the melting glaciers were responsible for most of the landforms that have an indirect relationship to glacial action. Most of these forms are mainly depositional and constructional, but meltwater trenches and terraces are at least partly erosional landforms. All the landforms in this category require the presence of ice in or near the area, but ice was not necessarily the primary mechanism responsible for determining the location, extent, or shape of the features.

Alluvial Plains

Areas underlain by gravel and sand are shown in yellow on Plate 1. Some of the gravel and sand is undoubtedly glacial outwash, but paleobotanical evidence suggests that a significant percentage of the material was deposited by rivers and streams that were fed by the increased precipitation that occurred in early postglacial time (9,000 to 12,500 B. P.) (Clayton, 1967, p. 33). The relative percentages of glacial outwash materials and postglacial alluvium are not known and, for purposes of discussion, the deposits will be referred to here simply as alluvial plains. The locations of these alluvial plains are shown on Plate 1.

The alluvial plains are relatively flat areas with local relief of 10 to 20 feet in a mile. Relief is greatest in areas where sand dunes have

developed. Channel and meander scars, none of which continue for appreciable distances, are common. They are most easily seen on air photos. Boulders are rare on the surface.

The alluvial plains of western and southern McLean County are underlain by fluvial sediments that are generally less than 10 feet thick, but in the Turtle Lake area over 100 feet of gravel and sand underlies the alluvial plain in places. It is unlikely, however, that the deeper fluvial sediments in these areas are related to the surficial alluvial plains; they probably were deposited during events that occurred prior to the deposition of the alluvial plains.

Collapsed Alluvial Plains

Hilly areas underlain by gravel and sand are collapsed alluvial plains. They consist of collapsed sediments of formerly flat alluvial plains that were deposited on stagnant glacial ice. When the stagnant ice melted, collapse of the overlying alluvium occurred. In general, these areas are characterized by hilly topography with abundant undrained depressions, faulted bedding and other collapse structures that can be observed in cuts, and disintegration trenches (Clayton, 1962, p. 42). Relief ranges up to about 75 feet in a square mile in the more rugged areas of collapsed alluvium.

In McLean County, numerous small linear bodies of collapsed alluvium occur on the Missouri Coteau (pl. 1). These are collapsed valley trains that were deposited in ice valleys on the stagnant glacier. A larger area of collapsed alluvium occurs in T. 147 N., Rs. 79 and 80 W. This area overlies a deep preglacial valley (pl. 3) that apparently became partly filled with stagnant ice that was later covered by fluvial deposits.

Kames

Kames are "mounds composed chiefly of gravel or sand, whose form has resulted from original deposition modified by any slumping incident to later melting of glacial ice against or upon which the deposit accumulated" (Holmes, 1947, p. 248). In areas of dead-ice moraine, kames should be restricted to conspicuous hills of sand and gravel that are more prominent than the surrounding hills. Numerous kames were mapped within the area of dead-ice moraine in McLean County, and a few were mapped in the area of ground moraine in the western part of the county (pl. 1). The most prominent kames are in western McLean County, where local relief is generally low. A good example is Paint Hill (SW $\frac{1}{4}$, Sec. 28, T. 149 N., R. 88 W.). This kame is about 50 feet high and about a quarter mile across at the base. It has few surface boulders, less than the surrounding area, and contains well sorted sand and gravel, some till, and a layer of silt.

Flat Lake Topography

In western McLean County, a relatively flat area of about 10 square miles occurs in T. 150 N., R. 86 W. This area, along with a similar slightly smaller area about 4 miles to the southwest, is underlain by banded silt and clay that is sandy in places. Relief over the two areas is generally less than 5 feet in a mile. These areas formed when the glacier that had deposited the Blue Mountain end moraine receded a short distance, and meltwater from the glacier was dammed between the ice and the end moraine.

Undulating Lake Topography

An undulating surface underlain by lake sediments, mainly silts and silty sands, that were deposited in a lake on stagnant or stagnating ice is lake topography that has collapsed. When the ice melted, the sediments became folded and contorted; and the surface, which had been flat, became undulating. Plate 1 shows several areas of collapsed lake deposits, all closely associated with dead-ice moraine. Some of the deposits are elevated slightly above the surrounding landscape and are rather flat. Apparently the lakes in which these silts and fine sands accumulated were surrounded by, but were not on, stagnant ice so that when the ice melted, surrounding areas collapsed and the lake topography was left intact. A few of these flat deposits have till rims at the edge indicating that debris slid from the adjacent ice into the lakes. On air photos the areas of collapsed lake topography are easily recognized as cultivated patches of ground of relatively low relief surrounded by uncultivated dead-ice moraine of higher relief.

Meltwater Trenches

Meltwater trenches are valleys that were cut by water flowing from melting ice sheets. Probably a significant amount of the water that flowed through these valleys was derived from local precipitation as well.

Many of the meltwater trenches of eastern McLean County, on the Missouri Coteau, are poorly defined due to collapse of the stagnant ice over and through which they were cut. Some of the areas shown as collapsed outwash topography on Plate 1 are confined in places to fairly well-defined valleys that are probably segments of meltwater trenches.

One of the largest meltwater trenches in the area of dead-ice moraine trends southward from the Ruso area into the outwash plain in T. 147 N., R. 81 W. This trench has a gravel floor and a single terrace level that grades to an upper, partially collapsed level within the outwash plain. Its floor grades to the lower, uncollapsed portion of the outwash plain.

Nearly paralleling this meltwater trench is another narrower but equally deep trench in Tps. 148 and 149 N., R. 80 W. Strawberry Lake, Long Lake, and Crooked Lake all occur within the trench. This trench was apparently cut while stagnant ice was still present. When the ice melted, the trench was blocked in places due to collapse as the original gradient was destroyed.

Further south is a deep, south-trending valley through which the Turtle Creek flows (pl. 1). The position of this trench, parallel to the edge of the late Wisconsinan dead-ice moraine, suggests it was initiated as an ice marginal feature. It begins near the town of Turtle Lake at the southern edge of the outwash plain referred to above. The trench is cut mainly in bedrock and has highly dissected walls with badland topography in places.

Another meltwater trench is the middle branch of Douglas Creek, which trends southward from the Ward County line to the Missouri River through R. 85 W. (pl. 1). This trench is also parallel to the edge of the late Wisconsinan dead-ice moraine and may have been initiated as an ice marginal feature. The trench is about a mile wide where it nears the Missouri River Valley (fig. 9) and about 150 feet deep. It is cut mainly in bedrock.

By far the largest meltwater trench in the area is the valley of the Missouri River. The Missouri River has already been discussed in the part of this paper dealing with drainage development. It was pointed out that some segments of the Missouri River trench are of preglacial origin; and others were cut by meltwater during glaciation, although the entire trench served as a meltwater route.

Erosion Surfaces

Relatively flat, pediment-like surfaces that are cut across both early Wisconsinan glacial drift and Tertiary bedrock occur in Tps. 144-145 N., Rs. 83-84 W. (fig. 11). The surfaces occur at elevations that range from 1,800 to 2,000 feet and they slope southeastward. In a few places, the erosion surfaces are covered by several feet of sandy gravel, but in most areas only a thin soil covers the bedrock or glacial drift. Boulders are common on the surfaces except in the gravel-covered areas. On air photos, the erosion surfaces look smooth and lack the morainic markings typical of the surrounding areas.

The erosion surfaces apparently were cut by water that flowed southeastward from the Riverdale area in late Wisconsinan time when east-trending drainage was blocked and the south-trending segment of the Missouri River had not yet been cut. By projecting the slopes on the erosion surfaces southeastward to the center of the Missouri River trench in the area west of Washburn, a figure of about 1,740 feet is arrived at for the elevations of the base of the Missouri River trench at




FIGURE 9. Middle branch of Douglas Creek meltwater trench in T. 148 N., R. 85 W. View north from state highway 37 in section 7.



FIGURE 10. Turtle Creek valley east of Washburn (T. 144 N., R. 81 W.). View is north over terraces along the meltwater trench.



FIGURE 11. View south over the flat profile of a remnant of an erosion surface (Secs. 25 and 26, T. 145 N., R. 84 W.). The low area in the foreground is the Missouri River floodplain.

the time the surfaces were cut. The present floor of the trench is at about 1,670 feet, so the valley today is about 70 feet deeper than it was in late Wisconsinan time.

Terraces

Many of the larger meltwater trenches have a single terrace level developed on the valley walls. In most cases these terraces are cut on either bedrock or till and have a covering of gravel that is generally less than 5 feet thick.

Except for very low local terrace levels that are generally less than 15 feet above river level, and considered to belong to the modern floodplain, three terrace levels can be recognized within the Missouri River trench. The lowest of these is about 40 to 50 feet above the river level at elevations that range from about 1,730 feet near Riverdale to less than 1,700 feet at the Burleigh County line. Upstream from Riverdale any low terraces that may have been present are now flooded by Lake Sakakawea. The lowest terrace level is underlain mainly by

gravel and is a high fill and cut terrace. It probably corresponds to terrace 4 in Burleigh County (Kume and Hansen, 1965, p. 8).

The middle terrace level is generally about 90 feet above the river level at elevations that range from about 1,790 in the Riverdale area down to about 1,735 near Washburn. This terrace level is cut across bedrock and early Wisconsinan till and has a gravel cover that is commonly less than 10 feet thick. The middle terrace is fairly continuous all the way from Riverdale to the Burleigh County line. Because this terrace level was cut after the modern route of the river was established in late Wisconsinan time, the level of the river has been lowered at least 90 feet since the route was established, presumably when the late Wisconsinan glacier reached its maximum extent.

The uppermost terrace level ranges from over 1,900 feet south of Garrison to about 1,770 feet near Washburn. This is a cut terrace with little alluvial material on the surface. It is apparently related to the cutting of the north-south segment of the trench south of Riverdale because pediments that were cut before the trench existed occur at about the same level.

NONGLACIAL FORMS

In McLean County only a few landforms are unrelated to glacial action and even these can be indirectly tied to the glaciations. The Missouri River floodplain, for example, is a modern feature but its distribution is controlled by its position in the Missouri River Trench, which is, in part at least, of glacial origin. Similarly, the location of the badlands is controlled by the fact that they occur primarily in the Missouri River Trench. This category of landforms includes features that were shaped or are being shaped by nonglacial processes. In contrast to the first two groups of landforms, which are mainly relict forms that have not changed significantly since they were deposited, the landforms in this category are still being formed today.

River Floodplains

Most of the modern streams in McLean County have floodplains that are only a few hundred feet wide and underlain by thin alluvium. The exception is the Missouri River floodplain which is as much as 3 to 4 miles wide. It consists of channel bars and meander bars that are continually being modified by present river action. Topography on the floodplain is generally smooth and level. Oxbow lakes, such as Painted Woods Lake in Secs. 9 and 16, T. 143 N., R. 80 W., occur in some of the old meander scars. The Missouri River floodplain is characterized by

heavily wooded areas in places. Cottonwood trees are particularly common.

Sloughs

Sloughs occur throughout McLean County, particularly in the areas of dead-ice moraine. Some of them become moderate-sized lakes during wet years, but normally they are areas of marshland. Most of the smaller sloughs occur in the bottoms of kettle depressions, but some, such as Weller Slough, (Secs. 1, 2, and 12, T. 145 N., R. 83 W.) and others in the Falkirk area, occur in broad depressions that may mark a buried valley.

Dunes

Sand dunes occur in Sec. 13, T. 145 N., R. 84 W., and Sec. 18, T. 145 N., R. 83 W. The source of the sand is the fluvial deposits on terraces of the Missouri River trench to the northwest. The dunes cover an area of about one square mile. They have local relief of about 10 feet between dune crests and adjacent depressions. A few blowouts occur in the dune area; but for the most part the dunes are stable with a grass and brush cover.

Stream-Eroded Topography

Areas of stream-eroded topography occur along the larger stream valleys in the area of the Napoleon drift where they coincide, approximately, with the areas of bedrock exposures (pl. 1). Only minor erosion has occurred on areas mantled by substantial thicknesses of glacial drift. The most extensive areas of stream-eroded topography occur within the Missouri River Trench. In Rs. 87-90 W. and Tps. 144-145 N., R. 84 W., the trench is characterized by badlands topography. These areas have slopes that range from 20° to 65° with locally high relief. Badlands are found on the Tongue River and Sentinel Butte Formations. A few buttes that are present in the badland areas are capped by glacial drift overlying a resistant sandstone or siliceous bedrock layer. Erosion of the badland topography has resulted in knife-like ridges with intervening arroyos and slopes that are almost devoid of vegetation.

GEOLOGIC HISTORY

Preglacial History

McLean County is located in the eastern part of the central Williston basin, the center of which is located southeast of Williston, North Dakota. Marine seas invaded this part of the Williston basin at least four times during Paleozoic time and deposited carbonates, sandstones, shales, and evaporites. During times the area was emergent, parts of these deposits were removed by erosion. The area was emergent during Early to Middle Ordovician, Middle Silurian to Middle Devonian, Late Mississippian to Early Pennsylvanian, and during the Triassic.

Marine deposition continued during the Mesozoic from Middle to Late Jurassic time at which time emergence with non-marine deposition occurred. This non-marine deposition continued into Cretaceous time until the Early Cretaceous seas invaded the area and thick accumulations of fine grained clastics were deposited. The last phase of marine Cretaceous deposition is represented by fine grained clastics of the Fox Hills Formation.

A change to non-marine deposition began in Late Cretaceous time with the deposition of the Hell Creek Formation. This continued into the Paleocene with the deposition of the Ludlow Formation. The last marine invasion of the area resulted in the deposition of the Cannonball Formation. This was followed by deposition of the non-marine Tongue River and Sentinel Butte Formations of Paleocene age. The Golden Valley Formation of Paleocene to Eocene age is present in Mercer County directly south of McLean County, and the presence of the Oligocene White River Formation in the Killdeer Mountains of Dunn County suggests that these two formations may have extended into McLean County, but if they were ever present, they were removed during the post-Oligocene erosional episode.

The post-Eocene Tertiary period was a time of development of an integrated drainage system over much of the area. During this time the predominant process was erosion of the then existing landscape. The topography of the area in early Pleistocene time was probably similar to the present topography in areas of thin glacial drift.

Topography on the Bedrock Surface

The topography of the bedrock surface under McLean County is shown on Plate 3. This generalized map is based largely on subsurface test hole data, and is therefore highly conjectural.

The preglacial surface, which was developed on the Tertiary Cannonball, Tongue River and Sentinel Butte Formations, was a moderately rolling plain dominated by three northeast-trending valleys. Elevations ranged from about 1,600 feet in the northeast corner of the county below the preglacial Missouri Escarpment to over 2,100 feet in the western part of the county. The largest river valley trended eastward along the southern edge of western McLean County through T. 147 N., Rs. 84-90 W. and continued eastward through T. 148 N., Rs. 79-83 W. (pl. 3). The part of the valley in the western half of the county coincides with the modern Missouri River valley, but the eastern part of the valley is buried beneath glacial drift.

The preglacial Knife River, which today joins the Missouri River about 10 miles south of Riverdale, continued northeastward to the valley described in the last paragraph in T. 148 N., R. 81 W. This northeast extension of the Knife River valley is today buried beneath glacial drift.

The third major preglacial valley shown on Plate 3 coincides with the Missouri River valley east of Washburn in T. 144 N., Rs. 82-84 W. and continues northeastward beneath the glacial drift to T. 148 N., R. 79 W., where it joins the first river valley.

In addition to the three preglacial valleys mentioned above, several rather narrow but deep meltwater trenches are shown on Plate 3. These trenches formed when drainage was diverted by the glaciers. In some places, the meltwater trenches may have been ice marginal.

Modern erosion has cut into the bedrock surface in some areas of thin glacial drift. This is most common along the edge of the Missouri River trench in western McLean County where rather extensive areas of Sentinel Butte Formation sands and silts are exposed at the surface.

Development of Drainage During the Quaternary

The map of the bedrock surface in McLean County (pl. 3) shows several rather deep valleys that cross the area. Some of these were in existence before the area was glaciated, and others were cut as glacial diversion trenches during the glacial epochs. In general, prior to

glaciation, the area was relatively flat with elevations on the uplands ranging from about 1,900 feet in the northeast to about 2,200 feet in the southwest. Valleys that trended generally east and northeast were cut about 300 feet into the upland surface and were about 4 to 6 miles wide.

PREGLACIAL DRAINAGE

Figure 12 shows a probable drainage pattern shortly prior to glaciation. These generally northeast-trending valleys contain fluvial sands and gravels of western derivation overlain by glacial deposits, which comprise most of the valley fill. The preglacial valleys were an important factor in establishing the modern course of the Missouri River, which marks the southwestern boundary of McLean County.

DRAINAGE DEVELOPMENT DURING GLACIATION

Those parts of the east-trending preglacial valleys that were not completely filled with glacial deposits again carried rivers during the glacial epochs. Due to increased precipitation during glacial times, these rivers were much larger than were the original preglacial rivers that had first cut the valleys and, as a result, trenches were cut in the wide valleys. Parts of the original preglacial valleys became blocked by glacial deposits and, in these areas, narrow diversion trenches were cut without regard to the regional slope. The modern Missouri River valley from Riverdale to a point about 12 miles south of Riverdale is an example of such a diversion trench. This relatively young segment of the Missouri River trench is much narrower than are east-trending segments of the trench that apparently carried preglacial drainage.

Many diversion trenches, now deeply buried in eastern and central North Dakota under glacial deposits, were cut as northeast-flowing streams became blocked by advancing glaciers and were diverted southeastward. As later glaciers advanced, they overrode ice-marginal trenches that had formed earlier. Eastern and northern areas of North Dakota are characterized by a myriad of channels now deeply buried beneath the drift and traceable only through test drilling. Examples of such buried diversion trenches in McLean County include the one that trends southeastward from T. 149 N., R. 90 W., to T. 147 N., R. 85 W. This trench is as much as 300 feet deep and has virtually no surface expression.

The vast array of buried valleys in McLean County as well as throughout eastern and northern North Dakota shows that drainages




FIGURE 12. Map showing major preglacial drainages and glacial diversion trenches in the McLean County area.

followed many routes during the various glaciations and the intervening interglacial stages. Since so many buried valleys have been found, it is difficult to pin down the drainage pattern that may have existed at any particular time prior to the present.

Some of the buried meltwater trenches in eastern North Dakota are both wide and deep, and it seems probable that they served as river valleys for long periods of time, perhaps for entire interglacial stages, carrying drainage from an area comparable in size to that drained by the modern Missouri River. Certainly, during the pre-Wisconsinan glaciations, drainage must have been southward around the ice sheets just as it was during the Wisconsinan. Some of the preglacial valleys were again used after each glaciation and some remained permanently blocked by glacial drift.

Regional evidence indicates that the modern Missouri River route did not exist downstream from a point near Riverdale in mid-Wisconsinan time. Instead, the main drainage flowed through the preglacial valley that trends eastward from Riverdale (fig. 12). South of Washburn, terraces of till-covered bedrock occur within the relatively narrow Missouri River Trench. The till is early Wisconsinan (?) in age, so this segment of the trench must have been cut as a glacial diversion feature sometime prior to the early Wisconsinan. Because the trench south of Washburn is relatively narrow, it must not have served as the main river route for a long period of time. It seems probable that this part of the trench was first cut during pre-Wisconsinan glaciations and then abandoned in favor of earlier, better established drainage further east when the ice receded.

The present route of the Missouri River from Riverdale southward was established during late Wisconsinan time. Large erosion surfaces were cut in the area southeast of Riverdale as the water flowed overland until a distinct trench could be cut. When a trench was established south of Riverdale, it was rapidly cut to a depth of about 300 feet. In contrast to the 4- to 6-mile-wide, till-draped valley upstream from Riverdale, the trench south of Riverdale is only about 1½ miles wide in places and is free of glacial drift.

Somewhere east of Washburn the preglacial drainage was again blocked by ice in late Wisconsinan time and was again diverted southward from near Washburn through the trench that had been cut earlier. The modern route of the Missouri River was thus finally established in late Wisconsinan time.

Smaller drainages such as Douglas Creek and Turtle Creek formed during late Wisconsinan time, although some of them tend to follow older, drift-lined valleys. Modern drainage over McLean County is poorly developed, particularly in the areas covered by late Wisconsinan drift.

Glacial History

INTRODUCTION

In the preceding pages, the Coleharbor Formation was named and the lithostratigraphic terminology necessary to characterize the formation was introduced. The value of lithostratigraphic units as descriptive units is due in part to the restriction that inferred geologic history plays no part in their definition. Each of the glacial drift units in McLean County is recognized on the basis of inferred geologic history or origin. For interpretive purposes, such inferential units are more useful than lithostratigraphic units, which are based on lithology.

Along the shore of Lake Sakakawea in McLean County, three glacial drifts are exposed in several places. The oldest of these, the Dead Man drift, (named for Dead Man Coulee where it was first observed) was deposited by pre-Wisconsinan ice. The till of the Dead Man drift is similar to till that was interpreted as pre-Wisconsinan found by the writer in eastern North Dakota in missile site excavations. Pre-Wisconsinan drift has also been tentatively identified by Clayton (1970) in Dunn County and in an area to the southeast of there (fig. 13). This, the Dunn drift, consists mainly of surface boulders. The Dead Man drift of McLean County is overlain by fluvial sediments and by a second drift, referred to here as the Mercer drift because it is widely exposed in Mercer County, to the southwest of McLean County. The Mercer drift is also considered to be of pre-Wisconsinan age. Leonard (1916) discussed pre-Wisconsinan till in western North Dakota. He apparently dealt collectively with the two lower tills exposed along Lake Sakakawea and the boulders discussed by Clayton.

Drift thought to be equivalent to the Napoleon drift of Logan and McIntosh Counties (Clayton, 1962) overlies the Mercer drift in the Lake Sakakawea exposures of McLean County. It was tentatively dated as early Wisconsinan (?) by Clayton. The Napoleon drift is exposed over a broad area of west central North Dakota (fig. 13). The Lostwood drift (named for Lostwood Lake in Mountrail County where it is well exposed) is the youngest glacial drift in McLean County. It covers much of eastern and northern North Dakota where it has been studied in considerable detail and broken down into minor components.

PRE-WISCONSINAN DRIFT

Dead Man Drift

The pre-Wisconsinan Dead Man drift is well-exposed along the shore of Lake Sakakawea in Sections 14, 15, 22, 23, 24 and 27, T. 147 N., R. 84 W. It was also observed in the NE $\frac{1}{4}$, SE $\frac{1}{4}$, Sec. 21, T. 144 N.,




FIGURE 13. Map of North Dakota showing maximum extent of pre-Wisconsinan, Early Wisconsinan and Late Wisconsinan glaciers. Areas covered by various aged drifts are shown. Shaded area is McLean County. (After Clayton, 1970).

R. 83 W., on the north side of the Missouri River trench. Its lower contact is submerged in most places, but it was observed to overlie bedrock in one exposure in Section 27, T. 147 N., R. 84 W.

The Dead Man drift consists of at least two distinct facies. The lowermost of these consists of 5 to 10 feet of iron-stained and poorly cemented gravel that grades to bedded silts in places and occurs in what appears to be a channel cut into the bedrock (Sec. 27, T. 147 N., R. 84 W.). It was not possible to determine the exact relationship between the silt deposit and the gravel. The gravel is a mixed assemblage of western lithologic types (quartz, chert, chalcedony, agate, and scoria) and glacially transported types (carbonates, granite, and varied metamorphics).

The upper facies of the Dead Man drift is mainly a silty to clayey till, 20 to 30 feet thick, that is oxidized in most places to a yellowish-buff color with irregular gray mottling. In a few deep wave-cuts the till is unoxidized at 15 to 20 feet below its upper surface. The oxidized-unoxidized boundary is very irregular. The till has conspicuous straight and irregular joints that have closely associated heavy accumulations of iron and manganese oxides.

The till of the Dead Man drift is harder and more compact than nearby overlying drifts. It is almost lithified in places and fresh faces broken from the till make single surfaces of a conglomerate. Most pebbles in the till are sound, but some of the igneous ones are weathered to ghosts and, uncommonly, the carbonate pebbles are rotted to shells. Commonly the fractures cut across and through the contained pebbles. Vertical cliffs are common in exposures along with associated horizontal ledges. Overlying drifts have been washed away by waves leaving either a clean surface on the more resistant Dead Man till or the gravel conglomerate of the overlying lower facies of the Mercer drift.

The Dead Man till contains some sandy pods up to about 2 feet across, but generally it is a uniform silty clay. It contains abundant chips, grains, and chunks of lignite up to several inches in diameter. Very large boulders of metamorphics and carbonates, 15 feet or more in diameter, are common in the till. Many of them show excellent grooves and striations. In places, large chunks of bedded silts occur within the till (fig. 15). The bedding is, without exception, distorted. These silt bodies were apparently picked up from a lake deposit and transported to their present location by the glacier that deposited the Dead Man drift. It is possible the silts are the same as the already mentioned ones of the lowermost facies of the Dead Man drift.

In some places the upper few feet of the Dead Man till looks like a paleosol with caliche accumulations on ped surfaces (fig. 17). The A horizon is absent and was apparently eroded away before the overlying



FIGURE 14. Nearly vertical exposure of boulder-clay (Dead Man till) showing wave-weathered joint faces.



FIGURE 15. Inclusion of banded silts (lake sediments) in boulder-clay (Dead Man till). Exposure is about 30 feet high.




FIGURE 16. "Cobbles" composed of boulder-clay (Dead Man till). Beaches along lake are covered with such rounded, water-worn pieces of boulder-clay.



FIGURE 17. Caliche accumulations near the top of the Dead Man till. This is apparently a paleosol.




FIGURE 18. Boulder-clay of unit 3 in description of type section of Coleharbor Formation (above shovel). Black pieces are lignite and areas of white are fresh, carbonate-covered joint faces. Light area on right below shovel is silt (unit 2) and area below and to left of shovel is a sandy gravel that was reworked when it was overridden by a glacier.

gravel was deposited. The upper surface of the Dead Man till is today being exhumed by wave action of Lake Sakakawea.

The Dead Man till may be equivalent to till that was found in eastern North Dakota in numerous Minuteman missile excavations west of the Pembina Escarpment. This till is commonly very hard, dense, bouldery, and highly jointed. It was deposited during a glaciation that is represented by about 200 feet of till in the Red River Valley (Bluemle, 1967). Knowledge of this glaciation is based mainly on a study of test hole data from the Red River Valley, particularly in Traill County, North Dakota (fig. 20).

Mercer Drift

The Mercer drift overlies the till of the Dead Man drift and consists of three facies, a lower fluvial deposit, a middle till deposit, and an upper fluvial deposit. The Mercer drift is exposed in most of the same areas as the Dead Man drift in McLean County, but it is widely



FIGURE 19. Contact between till and fluvial facies of the Mercer drift. Sand is loose and washes out easily leaving overhanging cliffs of Mercer till that tend to break off as large chunks and fall into the reservoir.




FIGURE 20. Comparison of glacial stratigraphy in McLean County with that in the Red River Valley.

exposed southwest of the Missouri River trench in Mercer County. In all exposures in McLean County, it is separated from the overlying early Wisconsinan drift by a sharp erosional contact. Its lower contact with the till of the Dead Man drift is also sharp in most places (fig. 6).

The lowermost facies of the Mercer drift is a discontinuous sandy gravel that forms a sharp contact with the underlying till. It averages about 3 to 4 feet thick. The gravel is highly iron-cemented in a few places, and pieces of it that have fallen into the reservoir have held together in chunks at least 10 feet long. In places, cross-bedded sand interfingers and lies above the gravel. In most places this sand is not even iron-stained. Such iron staining and cementation as is found in the gravels is apparently a function of local ground-water conditions. It has been suggested that iron staining and cementation are evidence that a deposit is unusually old, but the writer has observed similar conditions in gravels of late Wisconsinan age in Pembina County, North Dakota.

The iron-cemented gravels are apparently equivalent to those that have been informally called the Four Bears conglomerate (Clayton, in preparation). It is best exposed in the Sakakawea shore bluffs 1 mile northwest of the Four Bears Bridge in Mountrail County.

The middle facies of the Mercer drift is a till that is brownish gray, sandy, and pebbly with a few chips of lignite and pockets of reddish brown, finely divided limonite. It has strong vertical jointing that results in almost vertical cliffs (fig. 19). The Mercer till is about 25 feet thick; and, when viewed from a distance, it has a distinctive overall grayish aspect that makes it easy to distinguish from the underlying Dead Man drift and the overlying Napoleon drift. The till is partially oxidized throughout its thickness.

The till contains abundant inclusions of the underlying gravel and sand in some areas. As mentioned above, this gravel is iron-cemented in some locations where it is in place, but none of the inclusions that were found within the Mercer till were iron-cemented. This suggests that the iron cementing occurred after the gravel was buried beneath the Mercer till.

The gravel that overlies the Mercer till is up to 5 feet thick, sandy, crossbedded in some areas and has some iron staining near the base. It is discontinuous and looks like stream sediment that was deposited in a channel that was eroded into the surface of the till. If this is so, the gravel could be as old as the Mercer till or as young as the overlying Napoleon till.

PRE-WISCONSINAN GLACIAL HISTORY

Glaciers advanced in McLean County at least twice prior to the Wisconsin depositing the Dead Man and Mercer drifts. Stratigraphic relationships of the various facies of the Dead Man drift suggest that an advancing glacier caused a proglacial lake to form in the Riverdale area in an east-trending river valley. As the glacier continued to advance, probably southwestward, it overrode the silts that had been deposited in the lake and deposited the silty Dead Man till. It is not known how far the ice that deposited the Dead Man drift advanced. The boulders found at the surface in Dunn County may represent the remains of Dead Man drift that was deposited further west.

The Dead Man drift is unevenly oxidized throughout, and its upper surface is an erosion surface that appears to have been cut into a well-developed soil zone. The paleosol on top of the till of the Dead Man drift (fig. 17) suggests a significant interglacial interval; and, since the overlying Mercer drift is also pre-Wisconsinan, the Dead Man may be very old.

The gravel above the Dead Man till is probably only slightly older than the Mercer till that it lies beneath. It may be proglacial outwash material laid down in front of an advancing glacier. The absence of lake sediments suggests that no proglacial lake formed this time, perhaps indicating that the ice advanced from a northerly or northwesterly direction. However, this cannot be determined, as it is also possible that the east-trending preglacial drainage had been filled to such an extent during the first advance that a basin for a lake no longer existed.

Continued advance of the second glacier resulted in deposition of the Mercer till on the gravel. Again, it is not known how far this glacier advanced. The gravel that lies above the Mercer till is discontinuous and looks like a stream channel deposit. It may have been deposited any time after the Mercer till was deposited and before the Napoleon till was deposited.

EARLY WISCONSINAN DRIFT

Napoleon Drift

The early Wisconsin Napoleon drift covers a wide area of southern and western McLean County (fig. 21). It is bounded on the north and east by late Wisconsin Lostwood drift. The Napoleon drift averages about 15 feet thick over western McLean County to about 25 feet thick in the southern part of the county. It is probably considerably thicker in some of the buried trenches, but it is not known whether the materials it overlies in these trenches, usually sands or silts




FIGURE 21. Map of McLean County showing areas of various drifts.

of glacial origin, are part of the Napoleon drift or older drifts. Over most of western McLean County, the Napoleon drift has little or no morainic topography, but rather it is sheet moraine with almost completely integrated drainage. In the southern part of the county, subdued morainic topography is developed on the Napoleon drift in places, and only partially integrated drainage exists. These areas are subdued dead-ice moraine and ground moraine.

Till is the only lithologic type in nearly all the surface exposures of Napoleon drift in McLean County. Small pods of sand and gravel were seen in the till, but they are of very limited extent. In general, the till of the Napoleon drift is fairly silty to sandy but still rather cohesive. It is oxidized in all surface exposures to shades of yellowish brown to dusky yellow. The till is lignitic, although lignite particles tend to be smaller than do those of the Dead Man drift. Particles of limestone, shale, and granitic rocks are common. These range from highly weathered ghosts to sound pebbles and cobbles. In general, the Napoleon till is not significantly different than most North Dakota till. The thickness of oxidized sediments penetrated in some test holes spudded in Napoleon till is unusual. As much as 136 feet of continuous oxidized section consisting of till, sand, and silty clays occurred in test hole 2834 (Sec. 29, T. 148 N., R. 85 W.). In other holes, an interval of unoxidized sediment separates the surficial oxidized till from a second oxidized sediment, commonly till.

The Napoleon drift has not been distinguished lithologically in McLean County from the adjacent Lostwood drift. It is, however, easily distinguished geomorphically because the end moraines and the edge of the dead-ice moraine of the Lostwood drift overlie and truncate it. The resulting boundary on the Napoleon drift is readily apparent on air photos and in the field.

EARLY WISCONSINAN GLACIAL HISTORY

In early Wisconsinan time, perhaps 50,000 years ago, the Napoleon glacier advanced across North Dakota, until it covered all but the southwest corner of the state (fig. 13). While the Napoleon glacier was at its maximum extent, a deep diversion trench was cut at the edge of the ice through Dunn, Mercer, and Morton Counties by runoff from an area that was comparable in size to that now drained by the Missouri River. In McLean County, the Napoleon drift, where it is not covered by younger late Wisconsinan drift, consists of only a few tens of feet of till. For this reason, it seems unlikely the Napoleon ice covered the area for a very long period of time.

In southwestern McLean County, the Napoleon drift was deposited on a rolling surface that was mainly bedrock with only an occasional patch of pre-Wisconsinan drift in low areas. Most of the pre-Wisconsinan drift had been eroded away by the time Napoleon ice advanced over the area. In the eastern and northern parts of the county, however, considerable pre-Wisconsinan drift probably remained when the Napoleon drift was deposited. Numerous test holes in this area penetrated complex drift sections with several tills. Although it is impossible to correlate these buried tills with tills that are exposed further southwest, it is likely some of them are pre-Wisconsinan in age.

Figure 22 is necessarily generalized because the actual position of the Napoleon ice margin at any given time, as it receded from the area, is unknown. End moraines may have existed in the area, but they have since been overridden by late Wisconsinan ice and their positions are not known. It seems likely that by the time an approximation of the conditions portrayed in Figure 22 was reached, numerous meltwater diversion trenches, mainly in Mercer County, had been cut and, for the most part, abandoned. A proglacial lake formed about this time in the east-trending pre-glacial valley shown above and water overflowed southward, cutting an erosion surface in the area indicated. It seems likely that a rather thick layer of Napoleon drift was deposited on the Missouri Coteau. Wherever glaciers advanced over the Missouri Escarpment they probably picked up large volumes of material and subsequently deposited it on the Coteau. As the Napoleon glacier receded, it stagnated in places (fig. 22) leaving areas of drift-covered ice that later melted, resulting in dead-ice moraine.

LATE WISCONSINAN DRIFT

Lostwood Drift

Late Wisconsinan Lostwood drift covers the northern and eastern parts of McLean County (fig. 21). It is generally difficult to determine the thickness of the Lostwood drift in McLean County, because it is lithologically indistinguishable from underlying older drifts. Presumably it overlies early Wisconsinan Napoleon drift in most places, and in western McLean County the apparent contact is marked by an oxidized zone on top of the Napoleon drift in nine test holes. In these holes, the overlying Lostwood drift averaged 33 feet thick. In the eastern part of the county, very few test holes penetrated buried oxidized drift, so it was impossible to determine the thickness of the Lostwood drift there.

The Lostwood drift of McLean County can be subdivided into three areas with differing topographic expressions. In the western part of the county, the Blue Mountain end moraine marks the southern edge




FIGURE 22. Receding early Wisconsinan Napoleon ice.

of the Lostwood drift. Dead-ice moraine features of rather low relief are apparent on air photos in the area north of the Blue Mountain end moraine. Relief north of the end moraine is more subdued than is relief over most of the area of Lostwood drift in other parts of McLean County.

In eastern McLean County, the Lostwood drift is characterized by uneroded morainic topography with abundant dead-ice features. No prominent end moraines occur in eastern McLean County, but the boundary between Lostwood and Napoleon drift is easily identified by the change from integrated Napoleon sheet moraine to nonintegrated Lostwood dead-ice moraine. In extreme northeastern McLean County, the Lostwood drift is characterized by deeply eroded morainic topography on and adjacent to the Missouri Escarpment. This levels off to relatively uneroded morainic topography on the Drift Prairie to the northeast.

In most places the lithology of the Lostwood drift is essentially similar to that of the underlying Napoleon drift. The till is generally silty to sandy and slightly cohesive with abundant lignite chips. It is commonly oxidized in exposures to shades of yellowish brown. Most of the surficial gravel and sand in McLean County (pl. 1) is associated with the Lostwood drift. It is commonly a poorly sorted mixture of sand and angular to subangular gravel composed mainly of carbonates and chips of shale and lignite. Lake silts and sands that occur throughout the area are associated mainly with elevated lake plains formed only in the areas of dead-ice moraine. In extreme northeastern McLean County the Lostwood drift is unusual in that the tills contain high percentages of sand and gravel. In exposures, its till fabric is readily identified, but samples taken by hand auger commonly are indistinguishable from outwash deposits. In Sheridan County to the east, lineations can be seen on a gravel outwash surface indicating that the outwash was overridden by a glacier. This probably occurred in northeastern McLean County also, resulting in a till that is composed almost entirely of reworked fluvial sediments.

The late Wisconsinan age of the Lostwood drift has been verified by numerous radiocarbon dates. Most of the Lostwood drift in eastern McLean County was deposited from stagnant ice that may have taken 2,000 years to melt. Actively moving glaciers were probably present on the Missouri Coteau about 12,500 years ago. Radiocarbon dates of materials from Lostwood drift in North Dakota range from 9,000 B.P. (W-1019) to 11,070 B.P. (W-956) but these are from landforms deposited by stagnant ice. Materials taken from drift equivalent to Lostwood drift that have been dated in South Dakota range from 12,050 B.P. (W-1189) to 12,760 B.P. (Y-595).

Pettyjohn (1967) considers the Lostwood drift of western McLean and southwest Ward Counties (he refers to it as the Blue Mountain drift) to be of early Wisconsinan age. This is unlikely in view of the fact that stagnation features present on the Lostwood drift form an interlocking pattern with similar stagnation features all the way to the Missouri Escarpment. Such an interlocking pattern could result only if a continuous sheet of stagnant ice covered the Missouri Coteau and melted everywhere over that area at about the same time. Pettyjohn states that the Blue Mountain drift is oxidized to an average depth of 26 feet and that he has traced this oxidized zone nearly 40 miles north and more than 30 miles east of the Blue Mountain end moraine in Ward County. However, it is more likely that he was actually drilling into the Napoleon oxidized zone; certainly there is no reason to believe it is necessarily on the Blue Mountain drift. The unusually thick oxidized zone (44 feet) on the Blue Mountain drift of southwestern Ward County is probably a composite of oxidized Blue Mountain drift lying on oxidized Napoleon drift.

The total oxidized zone in areas covered by Lostwood drift in western McLean County averages about 35 feet thick. In view of the fact that the Lostwood drift itself averages only 33 feet thick in this area (based on 9 test holes), part of this oxidized interval must be in Napoleon drift. Because the two drifts are lithologically indistinguishable, it is impossible to determine how much of the oxidation has occurred since the Lostwood was deposited. In eastern McLean County, the oxidized zone on the Lostwood averages about 21 feet thick.

LATE WISCONSINAN GLACIAL HISTORY

In late Wisconsinan time, about 15,000 years ago, a glacier advanced across a large area of northern and eastern North Dakota. While it was at its maximum extent, two lobes of the glacier crossed McLean County. The westernmost of these (fig. 23) deposited the Blue Mountain end moraine (pl. 1) as it advanced southeastward. It apparently reached its maximum extent slightly earlier than did the lobe in eastern McLean County. Because the Blue Mountain end moraine was deposited against a bedrock high and the local slope was toward the ice, little outwash was deposited ahead of the end moraine. Instead, relatively narrow valleys were formed that carried the meltwater southward to a broad, east-trending valley that was later to carry the Missouri River. Figure 12 shows that much of the major pre-late Wisconsinan drainage in the McLean County area coincided with what is the modern Missouri River trench. During that part of




FIGURE 23. Deposition of the Blue Mountain end moraine.

early Wisconsinan time portrayed on Figure 23, the glacier diverted the major drainage southwestward in Mountrail County. The resulting diversion trench now carries the Missouri River, and the abandoned valley has become the Van Hook arm of Lake Sakakawea.

The glacier margin continued to advance in eastern McLean County until it reached the position shown on Figure 24. At the same time, the ice margin was receding from the Blue Mountain position to the west. The west-flowing part of the glacier overrode the Blue Mountain end moraine, but it seems unlikely that the resulting dead-ice moraine that truncates the end moraine is appreciably younger than is the end moraine. Disintegration features that are present on both ground moraine and low relief dead-ice moraine behind the Blue Mountain end moraine in Ward County form continuous chains with similar disintegration features to the northeast in high relief dead-ice moraine. This high relief dead-ice moraine was deposited from the northeast. This shows that a continuous area of stagnant ice existed throughout the area on both sides of the line of truncation. If there had been a significant time lapse between the deposition of the materials on either side of the line of truncation, the stagnant ice to the southwest would have melted, and no interfingering of disintegration features would have been possible.

When the ice margin receded from the Burnstad position, large areas of drift-covered stagnant ice were left behind. The active ice margin tended to become lobate as the glacier thinned and receded, and several small loop-shaped segments of recessional moraine were deposited, particularly in Sheridan County. Meltwater flowing from the glacier deposited thick sequences of gravel in eastern McLean County and southern Sheridan County. Much of this gravel was deposited largely on top of stagnant ice and later collapsed when the underlying ice melted. In the Turtle Lake area, broad areas of glacial outwash occur. The gravel occurs at two levels, the upper of which is slightly collapsed and the lower of which is flat. Apparently, stagnant ice was thin in the area. The first episode of gravel deposition was on the thin stagnant ice, but the second apparently occurred after the stagnant ice had melted.

By the time the ice margin had receded to the position shown on Figure 25, the modern course of the Missouri River was firmly established along the McLean County border. The diversion trenches southwest of the river were all abandoned and probably already partly filled with slopewash material.




FIGURE 24. Maximum extent of late Wisconsinan ice in eastern McLean County.




FIGURE 25. Withdrawal of the active ice margin from McLean County.

ECONOMIC GEOLOGY

Petroleum

Ten petroleum exploratory wells have been drilled in McLean County as of January 1, 1970. None of these has produced any oil and no oil shows were reported from any of the wells.

Lignite

Lignite occurs in the Tongue River and Sentinel Butte Formations of McLean County. Brant (1953) lists seven major lignite beds in the county but indicates some of these may correlate with one another. He estimates a total original reserve of 16,478,000,000 tons, but this figure is probably high. The total recoverable reserves are much lower than this, as strip mining techniques that are used to mine the lignite are limited to areas where overburden is less than about 100 feet thick.

Production of lignite in McLean County was 48,189 short tons in 1966 from two mines, the Underwood Coal Corporation, Inc., Underwood, and the B & W Coal Inc., Garrison. The total value of lignite production during the same period was \$176,129.00. Lignite produced in McLean County was valued at from \$3.50 to \$3.80 a ton.

Clinker

McLean County has a minor amount of clinker ("scoria"), a material formed when burning lignite beds bake the overlying sediments. Clinker is used locally for road surfacing material.

Riprap and Building Stone

Abundant glacial erratic boulders, found throughout McLean County, may be used as riprap. Riprap and building stone can also be obtained from the Tongue River and Sentinel Butte Formations, which contain resistant beds of sandstone.

REFERENCES CITED

- American Commission on Stratigraphic Nomenclature, 1961, Code of stratigraphic nomenclature: Am. Assoc. Petroleum Geologists Bull., v. 45, p. 645-665.
- Bluemle, J. P., 1967, Geology and ground water resources of Traill County, North Dakota, pt. 1, geology: North Dakota Geol. Survey Bull. 49, 34 p.
- Bluemle, J. P., 1970, Anomalous hills and associated depressions in central North Dakota: Geol. Soc. America (abstract), Program, 23rd annual meeting, Rocky Mountain Section, p. 325.
- Brant, R. A., 1953, Lignite resources of North Dakota: U. S. Geol. Survey Circ. 226, 78 p.
- Carlson, C. G., and Anderson, S. B., 1965, Sedimentary and tectonic history of North Dakota part of Williston basin: Am. Assoc. Petroleum Geologists Bull., v. 49, p. 1833-1846.
- Chamberlain, T. C., 1883, Terminal moraines of the second glacial epoch: U. S. Geol. Survey, 3rd. Ann. Rept., p. 291-402.
- Christiansen, E. A., 1967, A thin till in west-central Saskatchewan, Canada: Can. Jour. of Earth Science, v. 5, p. 329-336.
- Christiansen, E. A., 1968, Pleistocene stratigraphy of the Saskatoon area, Saskatchewan, Canada: Can. Jour. of Earth Science, v. 5, p. 1167-1173.
- Clayton, Lee, 1962, Glacial geology of Logan and McIntosh Counties, North Dakota: North Dakota Geol. Survey Bull. 37, 84 p.
- Clayton, Lee, 1966, Notes on Pleistocene stratigraphy of North Dakota: North Dakota Geol. Survey Rept. of Inv. 44, 25 p.
- Clayton, Lee, 1970, Geologic map of Dunn County, North Dakota: North Dakota Geol. Survey Misc. Map 11.
- Colton, R. B., Lemke, R. W., and Lindvall, R. M., 1963, Preliminary glacial map of North Dakota: U. S. Geol. Survey, Misc. Geol. Inv. Map I-331.
- Easton, G. B., 1966, Annual Report, July 1, 1964-June 30, 1966: North Dakota Coal Mine Inspection Department, 19 p.
- Holmes, C. D., 1947, Kames: Am. Jour. Sci., v. 245, p. 248.
- Krumbein, W. C., and Sloss, L. L., 1963, Stratigraphy and sedimentation: San Francisco, W. H. Freeman and Company, 660 p.
- Kume, Jack, 1964, Sheet moraine in Burleigh County, North Dakota: North Dakota Acad. Sci., v. 18, p. 162-166.
- Kume, Jack, and Hansen, D. E., 1965, Geology and ground water resources of Burleigh County, North Dakota, pt. 1, geology: North Dakota Geol. Survey Bull. 42, 111 p.

- Lemke, R. W., and Colton, R. B., 1958, Summary of Pleistocene geology of North Dakota, *in* Mid-Western Friends of the Pleistocene Guidebook, 9th. Ann. Field Conf.: North Dakota Geol. Survey Misc. Ser. 10, p. 41-57.
- Lemke, R. W., 1960, Geology of the Souris River area, North Dakota: U. S. Geol. Survey Prof. Paper 325, 138 p.
- Lemke, R. W., Laird, W. M., Tipton, M. J., and Lindvall, R. M., 1965, Quaternary geology of northern Great Plains *in* Wright, H. H., Jr., and Frey, D. G., The Quaternary of the United States: Princeton, Princeton University Press, p. 15-27.
- Leonard, A. G., 1916, The pre-Wisconsin drift of North Dakota: Jour. of Geology, v. 24, p. 521-532.
- Pettyjohn, W. A., 1967, Multiple drift sheets in southwestern Ward County, North Dakota, *in* Glacial geology of the Missouri Coteau and adjacent areas: North Dakota Geol. Survey Misc. Series 30, p. 123-129.
- Rau, J. L., Bakken, W. E., Chmelik, J. C., and Williams, B. J., 1962, Geology and ground water resources of Kidder County, North Dakota, Pt. I, Geology: North Dakota Geol. Survey Bull. 36, 70 p.
- Todd, J. E., 1896, The moraines of the Missouri Coteau and their attendant deposits: U. S. Geol. Survey Bull. 144, 71 p.
- Townsend, R. C., and Jenke, A. L., 1951, The problem of the origin of the Max moraine of North Dakota and Canada: Am. Jour. Sci., v. 249, p. 842-858.